

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁵ : C07K 15/04, A61K 39/29 G01N 33/569, C07K 7/10, 7/08 C07K 7/06</p>	<p>A2</p>	<p>(11) International Publication Number: WO 93/00365 (43) International Publication Date: 7 January 1993 (07.01.93)</p>
<p>(21) International Application Number: PCT/US92/05388 (22) International Filing Date: 24 June 1992 (24.06.92) (30) Priority data: 722,489 24 June 1991 (24.06.91) US (71) Applicant: CHIRON CORPORATION [US/US]; 4560 Horton Street, Emeryville, CA 94608 (US). (72) Inventors: CHIEN, David, Y. ; 1121 Douglas Court, Ala- mo, CA 94507 (US). RUTTER, William ; 80 Everson Street, San Francisco, CA 94131 (US). (74) Agents: BLACKBURN, Robert, P. et al.; Chiron Corpora- tion, Intellectual Property Department, 4560 Horton Street, Emeryville, CA 94608 (US).</p>		<p>(81) Designated States: AU, BG, CA, FI, HU, JP, KR, NO, PL, RO, RU, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LU, MC, NL, SE). Published <i>Without international search report and to be republished upon receipt of that report.</i></p>
<p>(54) Title: HEPATITIS C VIRUS (HCV) POLYPEPTIDES</p> <p>(57) Abstract</p> <p>HCV epitopes useful as immunogenic reagents, and corresponding antibodies and methods of use are disclosed.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FI	Finland	MI	Mali
AU	Australia	FR	France	MN	Mongolia
BB	Barbados	GA	Gabon	MR	Mauritania
BE	Belgium	GB	United Kingdom	MW	Malawi
BF	Burkina Faso	GN	Guinea	NI	Netherlands
BG	Bulgaria	GR	Greece	NO	Norway
BJ	Benin	HU	Hungary	PL	Poland
BR	Brazil	IE	Ireland	RO	Romania
CA	Canada	IT	Italy	RU	Russian Federation
CF	Central African Republic	JP	Japan	SD	Sudan
CG	Congo	KP	Democratic People's Republic of Korea	SE	Sweden
CH	Switzerland	KR	Republic of Korea	SN	Senegal
CI	Côte d'Ivoire	LI	Liechtenstein	SU	Soviet Union
CM	Cameroon	LK	Sri Lanka	TD	Chad
CS	Czechoslovakia	LU	Luxembourg	TC	Togo
DE	Germany	MC	Monaco	US	United States of America
DK	Denmark				

WO 93/00365

PCT/US92/05388

5

1

HEPATITIS C VIRUS (HCV) POLYPEPTIDES

10 Technical Field

The invention relates to materials and methodologies for managing the spread of hepatitis C virus (HCV) infection. More specifically, it relates to polypeptides useful as immunological reagents in the detection, prevention and treatment of HCV infections.

15

Background

HCV was first identified and characterized as a cause of non-A, non-B hepatitis (NANBH) by Houghton et al. This led to the disclosure of a number of general and specific polypeptides useful as immunological reagents.

20 *See, e.g.*, Houghton et al., EPO Pub. No. 318,216; Houghton et al., EPO Pub. No. 388,232; Choo et al., Science (1989) 244:359-362; Kuo et al., Science (1989) 244:362-364; Houghton et al., Hepatology (1991) 14:381-388. These publications provide the art with an extensive background on HCV generally, as well as the manufacture and uses of HCV polypeptide immunological reagents.

25 For brevity, therefore, the disclosure of these publications in particular are incorporated herein by reference.

Others have readily applied and extended the work of Houghton et al. *See, e.g.*, Highfield et al., UK Pat. App. 2,239,245 (The Wellcome Foundation Ltd.); Wang, EPO Pub. No. 442,394 (United Biomedical Inc.); Leung

30 et al., EPO Pub. No. 445,423 (Abbott Laboratories); Habits et al., EPO Pub. No. 451,891 (Akzo N.V.); Reyes et al., PCT Pub. No. WO 91/15516 (Genelabs Inc.);

WO 93/00365

PCT/US92/05388

2

Maki et al., EPO Pub. No. 468,657 (Tonen Corp.); and Kamada et al., EPO Pub. No. 469,348 (Shionogi Seiyaku K.K.).

Sensitive, specific methods for screening and identifying carriers of HCV and HCV-contaminated blood or blood products are an important advance in
5 medicine. Post-transfusion hepatitis (PTH) occurs in approximately 10% of transfused patients, and HCV has accounted for up to 90% of these cases. The major problem in this disease is the frequent progression to chronic liver damage (25-55%). Patient care as well as the prevention of transmission of HCV by
10 blood and blood products or by close personal contact require reliable diagnostic and prognostic tools, such as HCV polypeptides, to detect antibodies related to HCV. Such polypeptides are also useful as vaccines and immunotherapeutic therapeutic agents for the prevention and/or treatment of the disease.

Since HCV is a relatively new agent, a continuing need exists to define additional immunological reagents that will allow further study of the
15 clinical course of disease and the epidemiology of HCV in the population.

Disclosure of the Invention

The invention pertains to the characterization of new HCV epitopes. The characterization of these epitopes permits the manufacture of polypeptide
20 products which reacted immunologically with antibodies to HCV and/or generate anti-HCV antibody production in vivo. These polypeptide products are useful as standards or reagents in diagnostic tests and/or as components of vaccines. Antibodies, including for example both polyclonal and monoclonal, directed against HCV epitopes contained within these polypeptide sequences are also useful
25 reagents, for example, in diagnostic tests, as therapeutic agents, for screening of antiviral agents, and for the isolation of isolation/purification of HCV polypeptides or particles.

In its broadest sense, the present invention is directed to polypeptides containing the newly characterized HCV epitopes disclosed herein,
30 methods of manufacturing such polypeptides (e.g., recombinant and synthetic methods), methods of using such polypeptides (e.g., diagnostic, vaccine, and therapeutic), and articles of manufacture, compositions or formulations adapted to

WO 93/00365

PCT/US92/05388

3

such uses (e.g., polypeptides fixed to an immunoassay or other support, oral or injectable pharmaceutical compositions). Similarly, antibodies (polyclonal, monoclonal, or equivalents such as binding fragments, single-chain antigen-binding proteins, etc.) to the HCV epitopes disclosed herein are also included within the scope of the present invention, as well as methods of making such antibodies, methods of using such antibodies (e.g., diagnostic, vaccine, and therapeutic), and articles of manufacture, compositions or formulations adapted to such uses (e.g., antibodies fixed to an immunoassay or other support, oral or injectable pharmaceutical compositions).

Other aspects of the invention pertain to kits for analyzing samples for the presence of an HCV antigen comprising the above antibodies in a suitable container. Still other aspects of the invention pertain to kits for analyzing samples for the presence of an antibodies directed against an HCV antigen comprising a polypeptide as described above in a suitable container.

Still other aspects of the invention are: a method for producing a polypeptide containing the newly disclosed HCV epitopes comprising incubating host cells transformed with an expression vector containing a sequence encoding a polypeptide containing the HCV epitope under conditions which allow expression of said polypeptide; and a polypeptide containing such an HCV epitope produced by this method.

Immunoassays are also included in the invention. These include an immunoassay for detecting an HCV antigen comprising incubating a sample suspected of containing an HCV antigen with an antibody as described above under conditions which allow the formation of an antigen-antibody complex; and detecting an antigen-antibody complex containing the antibody. An immunoassay for detecting anti-HCV antibodies comprising incubating a sample suspected of containing anti-HCV antibodies with a polypeptide as described above, under conditions which allow the formation of an antibody-antigen complex; and detecting the antibody-antigen complex containing the polypeptide.

Also included in the invention are vaccines for treatment of HCV infection comprising an immunogenic peptide containing an HCV epitope described herein.

WO 93/00365

PCT/US92/05388

4

Yet another aspect of the invention is a method for producing antibodies to HCV comprising administering to an individual an isolated immunogenic polypeptide containing an HCV epitope described herein in an amount sufficient to produce an immune response.

5 The above aspects of the present invention are accomplished by the discovery of HCV epitopes of the formula

$$aa_x-aa_y$$

wherein aa denotes an amino acid;

x and y are integers such that $y-x \geq 6$;

10 aa_x-aa_y indicates a portion of the amino acid sequence of Figure 1;
and

x is selected from the group consisting of 23-34, 36, 66-79, 81-94, 96-98, 101-103, 186-189, 191, 206, 223, 232, 256, 286, 297-299, 321, 347, 357, 413, 414, 432, 465-471, 480-484, 501, 502, 521, 540-549, 579, 594-599, 601-
15 613, 641, 662-665, 685, 705, 706, 729, 782-789, 801, 851-855, 893, 916, 928, 946, 952-954, 1026, 1072, 1109, 1112-1117, 1218, 1240, 1280-1285, 1322, 1338, 1371, 1384, 1410, 1411, 1454, 1492, 1493, 1532-1535, 1560, 1561, 1566-1568, 1571-1577, 1601-1607, 1615-1620, 1655, 1695, 1710-1712, 1728, 1729, 1758-1762, 1781, 1808, 1821, 1851, 1880, 1908-1913, 1925, 1940-1948, 1951, 1966-
20 1969, 1999, 2001-2004, 2006-2014, 2024, 2048-2053, 2055-2057, 2071, 2088-2093, 2108, 2122-2148, 2165, 2187, 2226-2232, 2244-2249, 2267, 2281-2286, 2288, 2289, 2325-2327, 2346, 2347, 2349, 2382, 2401, 2417-2422, 2439-2444, 2446-2456, 2469, 2471-2476, 2495, 2533, 2534, 2573-2578, 2602-2604, 2606-2612, 2632-2638, 2660, 2676-2679, 2688-2693, 2707, 2721, 2757-2762, 2779,
25 2794, 2795, 2797-2799, 2801, 2802, 2817-2843, 2863-2867, 2878-2884, 2886-2895.

WO 93/00365

PCT/US92/05388

5

The above objects are also achieved using HCV epitopes of the formula

$$aa_x-aa_y$$

wherein aa denotes an amino acid;

5 x and y are integers such that $y-x \geq 6$;

aa_x-aa_y indicates a portion of the amino acid sequence of Figure 1;

and x is selected from the group consisting of 35 (where y is less than 45), 80 (where y is less than 90), 95 (where y is less than 110), 99 (where y is less than 120), 100 (where y is less than 150), 190 (where y is less than 210), 500 (where y is less than 550), 600 (where y is less than 625), 1260 (where y is less than 1280), 1569 (where y is less than 1931), 1570 (where y is less than 1590), 1694 (where y is less than 1735), 1949 (where y is less than 2124), 1950 (where y is less than 1985), 2000 (where y is less than 2050), 2005 (where y is less than 2025), 2054 (where y is less than 2223), 2250 (where y is less than 2330), 2287 (where y is less than 2385), 2290 (where y is less than 2310), 2345 (where y is less than 2375), 2348 (where y is less than 2464), 2445 (where y is less than 2475), 2470 (where y is less than 2490), 2605 (where y is less than 2620), 2780 (where y is less than 2830), 2796 (where y is less than 2886), 2800 (where y is less than 2850), and 2885 (where y is less than 2905).

20 In either of the above formula, x-y can less than or equal to 10, 20, 30, 40 or 50 in some embodiments of the invention.

Brief Description of the Drawings

Fig. 1 shows the polyprotein of the HCV prototype isolate HCV1.
25 Fig. 2 shows a composite cDNA sequence for HCV1.

Fig. 3 shows the nucleotide consensus sequence of human isolate 23, variant sequences are shown below the sequence line. The amino acids encoded in the consensus sequence are also shown.

Fig. 4 shows the nucleotide consensus sequence of human isolate 30 27, variant sequences are shown below the sequence line. The amino acids encoded in the consensus sequence are also shown.

WO 93/00365

PCT/US92/05388

Fig. 5 shows the aligned nucleotide sequences of human isolates 23 and 27 and of HCV1. Homologous sequences are indicated by the symbol (*). Non-homologous sequences are in small letters.

Fig. 6 shows the aligned amino acid sequences of human isolates 23 and 27 and of HCV1. Homologous sequences are indicated by the symbol (*). Non-homologous sequences are in small letters.

Fig. 7 shows a comparison of the composite aligned nucleotide sequences of isolates Thorn, EC1, HCT #18, and HCV1.

Fig. 8 shows a comparison of the nucleotide sequences of EC10 and a composite of the HCV1 sequence; the EC10 sequence is on the line above the dots, and the HCV1 sequence is on the line below the dots.

Fig. 9 shows a comparison of the amino acid sequences 117-308 (relative to HCV1) encoded in the "EnvL" regions of the consensus sequences of human isolates HCT #18, JH23, JH 27, Thorne, EC1, and of HCV1.

Fig. 10 shows a comparison of the amino acid sequences 330-360 (relative to HCV1) encoded in the "EnvR" regions of the consensus sequences of human isolates HCT #18, JH23, JH 27, Thorne, EC1, and of HCV1.

Modes for Carrying Out the Invention

Complete citations to publications referred to herein can be found in the "Background" or "Bibliography" sections.

I. Definitions

"Hepatitis C virus" or "HCV" refers to the art-recognized viral species of which pathogenic strains cause NANBH, and attenuated strains or defective interfering particles derived therefrom. See generally, publications cited in the section entitled "Background." The HCV genome is comprised of RNA. It is known that RNA containing viruses have relatively high rates of spontaneous mutation, *i.e.*, reportedly on the order of 10^{-3} to 10^{-4} per incorporated nucleotide (Fields & Knipe (1986)). Therefore, since heterogeneity and fluidity of genotype are inherent in RNA viruses, there are multiple strains/isolates, which may be virulent or avirulent, within the HCV species. The propagation, identification,

WO 93/00365

PCT/US92/05388

7

detection, and isolation of the various HCV strains or isolates has been well documented in the literature. Moreover, the disclosure herein allows the preparation of diagnostics and vaccines for the various strains/isolates, as well as compositions and methods that have utility in screening procedures for anti-viral agents for pharmacologic use, such as agents that inhibit replication of HCV.

Information on several different strains/isolates of HCV is disclosed herein, particularly strain or isolate CDC/HCV1 (also called HCV1). Information from one strain or isolate, such as a partial genomic or amino acid sequence, is sufficient to allow those skilled in the art using standard techniques to isolate new strains/isolates and to identify whether such new strains/isolates are HCV. For example, several different strains/isolates are described below. These strains, which were obtained from a number of human sera (and from different geographical areas), were isolated utilizing the information from the genomic sequence of HCV1.

The information provided herein is indicative that HCV may be distantly related to the flaviviridae. The Flavivirus family contains a large number of viruses which are small, enveloped pathogens of man. The morphology and composition of Flavivirus particles are known, and are discussed in Brinton (1986). Generally, with respect to morphology, Flaviviruses contain a central nucleocapsid surrounded by a lipid bilayer. Virions are spherical and have a diameter of about 40-50 nm. Their cores are about 25-30 nm in diameter. Along the outer surface of the virion envelope are projections that are about 5-10 nm long with terminal knobs about 2 nm in diameter. Typical examples of the family include Yellow Fever virus, West Nile virus, and Dengue Fever virus. They possess positive-stranded RNA genomes (~11,000 nucleotides) that are slightly larger than that of HCV and encode a polyprotein precursor of about 3500 amino acids. Individual viral proteins are cleaved from this precursor polypeptide.

The genomic structure and the nucleotide sequence of HCV genomic RNA has been deduced. The genome appears to be single-stranded RNA containing ~10,000 nucleotides. The genome is positive-stranded, and possesses a continuous, translational open reading frame (ORF) that encodes a polyprotein of about 3,000 amino acids. In the ORF, the structural protein(s) appear to be

WO 93/00365

PCT/US92/05388

8
encoded in approximately the first quarter of the N-terminus region, with the
majority of the polyprotein responsible for non-structural proteins. When
compared with all known viral sequences, small but significant co-linear
homologies are observed with the non-structural proteins of the flavivirus family,
5 and with the pestiviruses (which are now also considered to be part of the
Flavivirus family).

Based upon the putative amino acids encoded in the nucleotide
sequence of HCVI and other evidence, possible protein domains of the encoded
HCV polyprotein, as well as the approximate boundaries, are the following:

10

<u>Putative Domain</u>	<u>Approximate Boundary</u> <u>(amino acid nos.)</u>
15 C (nucleocapsid protein)	1-191
E ₁ (virion envelope protein)	192-383
20 E ₂ /NS1 (envelope?)	384-800
NS2 (unknown function)	800-1050
NS3 (protease?)	1050-1650
25 NS4 (unknown function)	1651-2100
NS5 (polymerase)	2100-3011(end)

WO 93/00365

-9-

PCT/US92/05388

These domains are, however, tentative. For example, the E1-NS2 border is probably in the 750-810 region, and NS3-NS4 border is about 1640-1650. There is also evidence that the 191 aa version of C is a precursor that is further processed (e.g., to about 170 aa in length), and that the NS2, NS4 and NS5 proteins are each further processed into two mature proteins.

Different strains, isolates or subtypes of HCV are expected to contain variations at the amino acid and nucleic acids compared with HCV1. Many isolates are expected to show much (*i.e.*, more than about 40%) homology in the total amino acid sequence compared with HCV1. However, it may also be found that there are other less homologous HCV isolates. These would be defined as HCV according to various criteria such as, for example, an ORF of approximately 9,000 nucleotides to approximately 12,000 nucleotides, encoding a polyprotein similar in size to that of HCV1, an encoded polyprotein of similar hydrophobic and/or antigenic character to that of HCV1, and the presence of co-linear peptide sequences that are conserved with HCV1. In addition, the genome would be a positive-stranded RNA.

HCV encodes at least one epitope which is immunologically identifiable with an epitope in the HCV1 polyprotein. The epitope is unique to HCV when compared to previously known Flaviviruses. The uniqueness of the epitope may be determined by its immunological reactivity with anti-HCV antibodies and lack of immunological reactivity with antibodies to known Flavivirus species. Methods for determining immunological reactivity are known in the art, for example, by radioimmunoassay, by ELISA assay, by hemagglutination, and several examples of suitable techniques for assays are provided herein. Alternatively, a comparison of the sequence of the HCV epitope to previously known sequences of members of the Flavivirus family can be used to evaluate "uniqueness."

In addition to the above, the following parameters of nucleic acid homology and amino acid homology are applicable, either alone or in combination, in identifying a strain/isolate as HCV. Since HCV strains and isolates are evolutionarily related, it is expected that the overall homology of the genomes at the nucleotide level may be about 10% or greater, probably will be

WO 93/00365

-10-

PCT/US92/05388

about 40% or greater, probably about 60% or greater, and even more probably about 80% or greater; and in addition that there will be corresponding contiguous sequences of at least about 13 nucleotides. It should be noted that variable and hypervariable regions within the HCV genome; therefore, the homology in these regions is expected to be significantly less than that in the overall genome. The correspondence between the putative HCV strain genomic sequence and, for example, the CDC/HCV1 cDNA sequence can be determined by techniques known in the art. For example, they can be determined by a direct comparison of the sequence information of the polynucleotide from the putative HCV, and the HCV cDNA sequence(s) described herein. For example, also, they can be determined by hybridization of the polynucleotides under conditions which form stable duplexes between homologous regions (for example, those which would be used prior to S₁ digestion), followed by digestion with single stranded specific nuclease(s), followed by size determination of the digested fragments.

Because of the evolutionary relationship of the strains or isolates of HCV, putative HCV strains or isolates are identifiable by their homology at the polypeptide level. Generally, HCV strains or isolates are expected to be at least 10% homologous, more than about 40% homologous, probably more than about 70% homologous, and even more probably more than about 80% homologous, and some may even be more than about 90% homologous at the polypeptide level. The techniques for determining amino acid sequence homology are known in the art. For example, the amino acid sequence may be determined directly and compared to the sequences provided herein. Alternatively the nucleotide sequence of the genomic material of the putative HCV may be determined (usually via a cDNA intermediate), the amino acid sequence encoded therein can be determined, and the corresponding regions compared.

As used herein, a polynucleotide "derived from" a designated sequence refers to a polynucleotide sequence which is comprised of a sequence of approximately at least about 6 nucleotides, preferably at least about 8 nucleotides, more preferably at least about 10-12 nucleotides, and even more preferably at least about 15-20 nucleotides corresponding to a region of the designated nucleotide sequence. "Corresponding" means homologous to or complementary to the designated sequence. Preferably, the sequence of the region from which the

WO 93/00365

-11-

PCT/US92/05388

polynucleotide is derived is homologous to or complementary to a sequence which is unique to an HCV genome. Whether or not a sequence is unique to the HCV genome can be determined by techniques known to those of skill in the art. For example, the sequence can be compared to sequences in databanks (as of the
5 priority date), *e.g.*, Genebank, to determine whether it is present in the uninfected host or other organisms. The sequence can also be compared to the known (as of the priority date) sequences of other viral agents, including those which are known to induce hepatitis, *e.g.*, HAV, HBV, and HDV, and to members of the Flaviviridae. The correspondence or non-correspondence of the derived sequence
10 to other sequences can also be determined by hybridization under the appropriate stringency conditions. Hybridization techniques for determining the complementarity of nucleic acid sequences are known in the art. See, for example, Maniatis *et al.* (1982). In addition, mismatches of duplex polynucleotides formed by hybridization can be determined by known techniques,
15 including for example, digestion with a nuclease such as S1 that specifically digests single-stranded areas in duplex polynucleotides. Regions from which typical DNA sequences may be "derived" include but are not limited to, for example, regions encoding specific epitopes, as well as non-transcribed and/or non-translated regions.

20 The derived polynucleotide is not necessarily physically derived from the nucleotide sequence shown, but may be generated in any manner, including for example, chemical synthesis or DNA replication or reverse transcription or transcription. In addition, combinations of regions corresponding to that of the designated sequence may be modified in ways known in the art to be
25 consistent with an intended use.

Similarly, a polypeptide or amino acid sequence "derived from" a designated amino acid or nucleic acid sequence refers to a polypeptide having an amino acid sequence identical to that of a polypeptide encoded in the sequence, or a portion thereof wherein the portion consists of at least 3-5 amino acids, and
30 more preferably at least 8-10 amino acids, and even more preferably at least 11-15 amino acids, or which is immunologically identifiable with a polypeptide encoded in the sequence. This terminology also includes a polypeptide expressed from a designated nucleic acid sequence.

WO 93/00365

-12-

PCT/US92/05388

A recombinant or derived polypeptide is not necessarily translated from a designated nucleic acid sequence; it may be generated in any manner, including for example, chemical synthesis, or expression of a recombinant expression system, or isolation from HCV, including mutated HCV. A recombinant or derived polypeptide may include one or more analogs of amino acids or unnatural amino acids in its sequence. Methods of inserting analogs of amino acids into a sequence are known in the art. It also may include one or more labels, which are known to those of skill in the art.

The term "recombinant polynucleotide" as used herein intends a polynucleotide of genomic, cDNA, semisynthetic, or synthetic origin which, by virtue of its origin or manipulation: (1) is not associated with all or a portion of a polynucleotide with which it is associated in nature, (2) is linked to a polynucleotide other than that to which it is linked in nature, or (3) does not occur in nature.

The term "polynucleotide" as used herein refers to a polymeric form of nucleotides of any length, either ribonucleotides or deoxyribonucleotides. This term refers only to the primary structure of the molecule. Thus, this term includes double- and single-stranded DNA and RNA. It also includes known types of modifications, for example, labels which are known in the art, methylation, "caps", substitution of one or more of the naturally occurring nucleotides with an analog, internucleotide modifications such as, for example, those with uncharged linkages (*e.g.*, methyl phosphonates, phosphotriesters, phosphoramidates, carbamates, *etc.*) and with charged linkages (*e.g.*, phosphorothioates, phosphorodithioates, *etc.*), those containing pendant moieties, such as, for example proteins (including for *e.g.*, nucleases, toxins, antibodies, signal peptides, poly-L-lysine, *etc.*), those with intercalators (*e.g.*, acridine, psoralen, *etc.*), those containing chelators (*e.g.*, metals, radioactive metals, boron, oxidative metals, *etc.*), those containing alkylators, those with modified linkages (*e.g.*, alpha anomeric nucleic acids, *etc.*), as well as unmodified forms of the polynucleotide.

A "purified" polypeptide refers to the polypeptide being in a state that is substantially free of other polypeptides, *i.e.*, in a composition that contains a minimum of about 50% by weight (desired polypeptide/total polypeptide in composition), preferably a minimum of about 70%, and even more preferably a

WO 93/00365

-13-

PCT/US92/05388

minimum of about 90% of the desired polypeptide, without regard to non-proteinaceous materials in the composition. Techniques for purifying viral polypeptides are known in the art. Purified antibodies are similarly defined.

"Recombinant host cells", "host cells", "cells", "cell lines", "cell cultures", and other such terms denoting microorganisms or higher eukaryotic cell lines cultured as unicellular entities refer to cells which can be, or have been, used as recipients for recombinant vector or other transfer DNA, and include the progeny of the original cell which has been transfected. It is understood that the progeny of a single parental cell may not necessarily be completely identical in morphology or in genomic or total DNA complement as the original parent, due to natural, accidental, or deliberate mutation.

A "replicon" is any genetic element, *e.g.*, a plasmid, a chromosome, a virus, a cosmid, *etc.* that behaves as an autonomous unit of polynucleotide replication within a cell; *i.e.*, capable of replication under its own control.

A "vector" is a replicon in which another polynucleotide segment is attached, so as to bring about the replication and/or expression of the attached segment.

"Control sequence" refers to polynucleotide sequences which are necessary to effect the expression of coding sequences to which they are ligated. The nature of such control sequences differs depending upon the host organism; in prokaryotes, such control sequences generally include promoter, ribosomal binding site, and terminators; in eukaryotes, generally, such control sequences include promoters, terminators and, in some instances, enhancers. The term "control sequences" is intended to include, at a minimum, all components whose presence is necessary for expression, and may also include additional components whose presence is advantageous, for example, leader sequences.

"Operably linked" refers to a juxtaposition wherein the components so described are in a relationship permitting them to function in their intended manner. A control sequence "operably linked" to a coding sequence is ligated in such a way that expression of the coding sequence is achieved under conditions compatible with the control sequences.

WO 93/00365

-14-

PCT/US92/05388

An "open reading frame" (ORF) is a region of a polynucleotide sequence which encodes a polypeptide; this region may represent a portion of a coding sequence or a total coding sequence.

5 A "coding sequence" is a polynucleotide sequence which is transcribed into mRNA and/or translated into a polypeptide when placed under the control of appropriate regulatory sequences. The boundaries of the coding sequence are determined by a translation start codon at the 5'-terminus and a translation stop codon at the 3'-terminus. A coding sequence can include, but is not limited to mRNA, cDNA, and recombinant polynucleotide sequences.

10 "Immunologically identifiable with/as" refers to the presence of epitope(s) and polypeptides(s) which are also present in the designated polypeptide(s), usually HCV proteins. Immunological identity may be determined by antibody binding and/or competition in binding; these techniques are known to those of average skill in the art.

15 As used herein, "epitope" refers to an antigenic determinant of a polypeptide. An epitope could comprise 3 or more amino acids that defines the binding site of an antibody. Generally an epitope consists of at least 5 amino acids, and sometimes consists of at least 8 amino acids. Methods of epitope mapping are known in the art.

20 A polypeptide is "immunologically reactive" with an antibody when it binds to an antibody due to antibody recognition of a specific epitope contained within the polypeptide. Immunological reactivity may be determined by antibody binding, more particularly by the kinetics of antibody binding, and/or by competition in binding using as competitor(s) a known polypeptide(s) containing an epitope
25 against which the antibody is directed. The techniques for determining whether a polypeptide is immunologically reactive with an antibody are known in the art.

30 As used herein, the term "antibody" refers to a polypeptide or group of polypeptides which are comprised of at least one antibody combining site. An "antibody combining site" or "binding domain" is formed from the folding of variable domains of an antibody molecule(s) to form three-dimensional binding spaces with an internal surface shape and charge distribution complementary to the features of an epitope of an antigen, which allows an immunological reaction with the antigen. An antibody combining site may be formed from a heavy and/or a

WO 93/00365

-15-

PCT/US92/05388

light chain domain (V_H and V_L , respectively), which form hypervariable loops which contribute to antigen binding. The term "antibody" includes, for example, vertebrate antibodies, hybrid antibodies, chimeric antibodies, altered antibodies, univalent antibodies, the Fab proteins, and single domain antibodies.

5 As used herein, a "single domain antibody" (dAb) is an antibody which is comprised of an V_H domain, which reacts immunologically with a designated antigen. A dAb does not contain a V_L domain, but may contain other antigen binding domains known to exist in antibodies, for example, the kappa and lambda domains. Methods for preparing dAbs are known in the art. See, for
10 example, Ward *et al.* (1989).

Antibodies may also be comprised of V_H and V_L domains, as well as other known antigen binding domains. Examples of these types of antibodies and methods for their preparation are known in the art (see, *e.g.*, U.S. Patent No. 4,816,467, which is incorporated herein by reference), and include the following.
15 For example, "vertebrate antibodies" refers to antibodies which are tetramers or aggregates thereof, comprising light and heavy chains which are usually aggregated in a "Y" configuration and which may or may not have covalent linkages between the chains. In vertebrate antibodies, the amino acid sequences of all the chains of a particular antibody are homologous with the chains found in one
20 antibody produced by the lymphocyte which produces that antibody *in situ*, or *in vitro* (for example, in hybridomas). Vertebrate antibodies typically include native antibodies, for example, purified polyclonal antibodies and monoclonal antibodies. Examples of the methods for the preparation of these antibodies are described *infra*.

25 "Hybrid antibodies" are antibodies wherein one pair of heavy and light chains is homologous to those in a first antibody, while the other pair of heavy and light chains is homologous to those in a different second antibody. Typically, each of these two pairs will bind different epitopes, particularly on different antigens. This results in the property of "divalence", *i.e.*, the ability to
30 bind two antigens simultaneously. Such hybrids may also be formed using chimeric chains, as set forth below.

"Chimeric antibodies", are antibodies in which the heavy and/or light chains are fusion proteins. Typically the constant domain of the chains is

WO 93/00365

-16-

PCT/US92/05388

from one particular species and/or class, and the variable domains are from a different species and/or class. Also included is any antibody in which either or both of the heavy or light chains are composed of combinations of sequences mimicking the sequences in antibodies of different sources, whether these sources
5 be differing classes, or different species of origin, and whether or not the fusion point is at the variable/constant boundary. Thus, it is possible to produce antibodies in which neither the constant nor the variable region mimic known antibody sequences. It then becomes possible, for example, to construct antibodies whose variable region has a higher specific affinity for a particular antigen, or
10 whose constant region can elicit enhanced complement fixation, or to make other improvements in properties possessed by a particular constant region.

Another example is "altered antibodies", which refers to antibodies in which the naturally occurring amino acid sequence in a vertebrate antibody has been varied. Utilizing recombinant DNA techniques, antibodies can be redesigned
15 to obtain desired characteristics. The possible variations are many, and range from the changing of one or more amino acids to the complete redesign of a region, for example, the constant region. Changes in the constant region, in general, to attain desired cellular process characteristics, *e.g.*, changes in complement fixation, interaction with membranes, and other effector functions.
20 Changes in the variable region may be made to alter antigen binding characteristics. The antibody may also be engineered to aid the specific delivery of a molecule or substance to a specific cell or tissue site. The desired alterations may be made by known techniques in molecular biology, *e.g.*, recombinant techniques, site directed mutagenesis, *etc.*

25 Yet another example are "univalent antibodies", which are aggregates comprised of a heavy chain/light chain dimer bound to the Fc (*i.e.*, constant) region of a second heavy chain. This type of antibody escapes antigenic modulation. See, *e.g.*, Glennie *et al.* (1982).

Included also within the definition of antibodies are "Fab" fragments
30 of antibodies. The "Fab" region refers to those portions of the heavy and light chains which are roughly equivalent, or analogous, to the sequences which comprise the branch portion of the heavy and light chains, and which have been shown to exhibit immunological binding to a specified antigen, but which lack the

WO 93/00365

-17-

PCT/US92/05388

effector Fc portion. "Fab" includes aggregates of one heavy and one light chain (commonly known as Fab'), as well as tetramers containing the 2H and 2L chains (referred to as F(ab)₂), which are capable of selectively reacting with a designated antigen or antigen family. "Fab" antibodies may be divided into subsets analogous to those described above, i.e., "vertebrate Fab", "hybrid Fab", "chimeric Fab", and "altered Fab". Methods of producing "Fab" fragments of antibodies are known within the art and include, for example, proteolysis, and synthesis by recombinant techniques.

Also included in the term "antibodies" are single-chain antigen-binding (SCA) proteins, such as the type described in the article co-authored by Schlom, J. in the June 15, 1992 issue of Cancer Research (as well as articles cited therein).

As used herein, the term "immunogenic polypeptide" is a polypeptide that elicits a cellular and/or humoral immune response, whether alone or linked to a carrier in the presence or absence of an adjuvant.

The term "polypeptide" refers to a polymer of amino acids and does not refer to a specific length of the product; thus, peptides, oligopeptides, and proteins are included within the definition of polypeptide. This term also does not refer to or exclude post-expression modifications of the polypeptide, for example, glycosylation, acetylation, phosphorylation and the like. Included within the definition are, for example, polypeptides containing one or more analogs of an amino acid (including, for example, unnatural amino acids, *etc.*), polypeptides with substituted linkages, as well as other modifications known in the art, both naturally occurring and non-naturally occurring.

"Transformation", as used herein, refers to the insertion of an exogenous polynucleotide into a host cell, irrespective of the method used for the insertion, for example, direct uptake, transduction, f-mating or electroporation. The exogenous polynucleotide may be maintained as a non-integrated vector, for example, a plasmid, or alternatively, may be integrated into the host genome.

"Treatment" as used herein refers to prophylaxis and/or therapy.

An "individual", as used herein, refers to vertebrates, particularly members of the mammalian species, and includes but is not limited to animals

WO 93/00365

-18-

PCT/US92/05388

(e.g., dogs, cats, cattle, swine, sheep, goat, rabbits, mice, rats, guinea pigs, etc.), and primates, including monkeys, chimps, baboons and humans.

As used herein, the "sense strand" of a nucleic acid contains the sequence that has sequence homology to that of mRNA. The "anti-sense strand" contains a sequence which is complementary to that of the "sense strand".

As used herein, a "positive stranded genome" of a virus is one in which the genome, whether RNA or DNA, is single-stranded and which encodes a viral polypeptide(s). Examples of positive stranded RNA viruses include Togaviridae, Coronaviridae, Retroviridae, Picornaviridae, and Caliciviridae. Included also, are the Flaviviridae, which were formerly classified as Togaviridae. See Fields & Knipe (1986).

As used herein, "antibody containing body component" refers to a component of an individual's body which is a source of the antibodies of interest. Antibody containing body components are known in the art, and include but are not limited to, for example, plasma, serum, spinal fluid, lymph fluid, the external sections of the respiratory, intestinal, and genitourinary tracts, tears, saliva, milk, white blood cells, and myelomas.

As used herein, a "biological sample" refers to a sample of tissue or fluid isolated from an individual, including but not limited to, for example, plasma, serum, spinal fluid, lymph fluid, the external sections of the skin, respiratory, intestinal, and genitourinary tracts, tears, saliva, milk, blood cells, tumors, organs, and also samples of *in vitro* cell culture constituents (including but not limited to conditioned medium resulting from the growth of cells in cell culture medium, putatively virally infected cells, recombinant cells, and cell components).

25

II. Description of the Invention

The practice of the present invention will employ, unless otherwise indicated, conventional techniques of molecular biology, microbiology, recombinant DNA, and immunology, which are within the skill of the art. Such techniques are explained fully in the literature. See e.g., Maniatis, Fitch & Sambrook, "Molecular Cloning; A Laboratory Manual" (1982); "DNA Cloning, Volumes I and II" (D.N Glover ed. 1985); "Oligonucleotide Synthesis" (M.J. Gait ed, 1984); "Nucleic Acid Hybridization" (B.D. Hames & S.J. Higgins eds. 1984);

30

WO 93/00365

-19-

PCT/US92/05388

"Transcription and Translation" (B.D. Hames & S.J. Higgins eds. 1984);
"Animal Cell Culture" (R.I. Freshney ed. 1986); "Immobilized Cells And
Enzymes" (IRL Press, 1986); B. Perbal, "A Practical Guide To Molecular
Cloning" (1984); the series, "Methods in Enzymology" (Academic Press, Inc.);
5 "Gene Transfer Vectors For Mammalian Cells" (J.H. Miller and M.P. Calos eds.
1987, Cold Spring Harbor Laboratory), Meth Enzymol Vol. 154 and Vol. 155
(Wu and Grossman, and Wu, eds., respectively), Mayer and Walker, eds. (1987),
"Immunochemical Methods In Cell And Molecular Biology" (Academic Press,
London); Scopes, (1987) "Protein Purification: Principles and Practice", Second
10 Edition (Springer-Verlag, N.Y.); and "Handbook of Experimental Immunology",
Volumes I-IV (D.M. Weir and C. C. Blackwell eds 1986). All patents, patent
applications, and publications mentioned herein, both *supra* and *infra*, are hereby
incorporated herein by reference.

15 II.A. Truncated HCV Polypeptides

The useful materials and processes of the present invention are
made possible by the identification below of new HCV epitopes. The knowledge
of these epitopes (or antigenic regions) allows for construction of polypeptides
containing truncated HCV sequences which can be used as immunological
20 reagents.

Truncated HCV amino acid sequences encoding at least one viral
epitope are useful immunological reagents. For example, polypeptides comprising
such truncated sequences can be used as reagents in an immunoassay. These
polypeptides also are candidate subunit antigens in compositions for antiserum
25 production or vaccines. While these truncated sequences can be produced by
various known treatments of native viral protein, it is generally preferred to make
synthetic or recombinant polypeptides comprising an HCV sequence. Polypeptides
comprising these truncated HCV sequences can be made up entirely of HCV
sequences (one or more epitopes, either contiguous or noncontiguous), or HCV
30 sequences and heterologous sequences in a fusion protein. Useful heterologous
sequences include sequences that provide for secretion from a recombinant host,
enhance the immunological reactivity of the HCV epitope(s), or facilitate the
coupling of the polypeptide to an immunoassay support or a vaccine carrier. See,

WO 93/00365

-20-

PCT/US92/05388

e.g., EPO Pub. No. 116,201; U.S. Pat. No. 4,722,840; EPO Pub. No. 259,149; U.S. Pat. No. 4,629,783, the disclosures of which are incorporated herein by reference.

The size of polypeptides comprising the truncated HCV sequences can vary widely, the minimum size being a sequence of sufficient size to provide an HCV epitope, while the maximum size is not critical. For convenience, the maximum size usually is not substantially greater than that required to provide the desired HCV epitopes and function(s) of the heterologous sequence, if any. Typically, the truncated HCV amino acid sequence will range from about 5 (or 8) to about 100 amino acids in length. More typically, however, the HCV sequence will be a maximum of about 50 (or 40) amino acids in length, and sometimes a maximum of about 20, 25 or 30 amino acids. It is usually desirable to select HCV sequences of at least about 8, 10, 12 or 15 amino acids.

Examples of truncated HCV amino acid sequences (octamers) that are useful as described herein are set forth below in the examples. It is to be understood that these peptides do not necessarily precisely map one epitope. Non-immunogenic portions of the sequence can be defined using conventional techniques and deleted from the described sequences. Further, additional truncated HCV amino acid sequences that comprise an epitope or are immunogenic can be identified as described herein.

Polypeptide products containing the truncated HCV amino acid sequences disclosed below can be prepared as discrete peptides or incorporated into a larger polypeptide, and may find use as described herein. In preferred applications, truncated sequences from the E1 and/or E2 domains have applications in vaccine and therapeutic products. While generally any of the domains can have some diagnostic utility, C, NS3, NS4 and NS5 are particularly preferred, with combinations of C epitopes with epitopes from one or more of the NS3, NS4 or NS5 domains being particularly preferred.

30 II.B. Preparation of Polypeptides

The availability of DNA sequences encoding HCV amino acid sequences permits the construction of expression vectors encoding antigenically active regions of the polypeptide (See, *e.g.*, Fig. 2). These antigenically active

WO 93/00365

-21-

PCT/US92/05388

regions may be derived from coat or envelope antigens or from core antigens, or from antigens which are non-structural including, for example, polynucleotide binding proteins, polynucleotide polymerase(s), and other viral proteins required for the replication and/or assembly of the virus particle. Fragments encoding the
5 desired polypeptides are derived, for example, from viral cDNA clones using conventional restriction digestion or by synthetic methods, and are ligated into vectors which may, for example, contain portions of fusion sequences such as β -Galactosidase or superoxide dismutase (SOD), preferably SOD. Methods and vectors which are useful for the production of polypeptides which contain fusion
10 sequences of SOD are described in European Patent Office Publication number 0196056, published October 1, 1986. Vectors encoding fusion polypeptides of SOD and HCV polypeptides, *i.e.*, NANB₅₋₁₋₁, NANB₈₁, and C100-3, which is encoded in a composite of HCV cDNAs, are described in Sections IV.B.1, IV.B.2, and IV.B.4, respectively. Any desired portion of the HCV cDNA
15 containing an open reading frame (or a synthetic version thereof), can be used to express a recombinant polypeptide, such as a mature or fusion protein.

Alternatively, a polypeptide contain HCV epitopes can be provided by chemical synthesis using standard techniques based on the amino acid sequence of the figures and the examples.

20 The DNA encoding the desired polypeptide, whether in fused or unfused form, and whether or not containing a signal sequence to permit secretion, may be ligated into expression vectors suitable for any convenient host. Both eukaryotic and prokaryotic host systems are presently used in forming recombinant polypeptides, and host cell lines is given in EPO Pub. Nos. 318,216. The
25 polypeptide is then isolated from lysed cells or from the culture medium and purified to the extent needed for its intended use. Purification may be by techniques known in the art, for example, differential extraction, salt fractionation, chromatography on ion exchange resins, affinity chromatography, centrifugation, and the like. See, for example, Methods in Enzymology for a variety of methods
30 for purifying proteins. Such polypeptides can be used as diagnostics, or those which give rise to neutralizing antibodies may be formulated into vaccines. Anti-bodies raised against these polypeptides can also be used as diagnostics, or for

WO 93/00365

-22-

PCT/US92/05388

passive immunotherapy. In addition, as discussed below, antibodies to these polypeptides are useful for example in isolating and identifying HCV particles.

The HCV polypeptides may also be isolated from HCV virions and truncated (if not already). The virions may be grown in HCV infected cells in
5 tissue culture, or in an infected host.

II.C. Preparation of Antigenic Polypeptides and Conjugation with Carrier

An antigenic region of a polypeptide is generally relatively small—typically 8 to 10 amino acids or less in length. Fragments of as few as 5
10 amino acids may characterize an antigenic region. These segments may correspond to regions of HCV antigen. Accordingly, using the cDNAs of HCV as a basis, DNAs encoding short segments of HCV polypeptides can be expressed recombinantly either as fusion proteins, or as isolated polypeptides. In addition, short amino acid sequences can be conveniently obtained by chemical synthesis.
15 In instances wherein the synthesized polypeptide is correctly configured so as to provide the correct epitope, but is too small to be immunogenic, the polypeptide may be linked to a suitable carrier.

A number of techniques for obtaining such linkage are known in the art, including the formation of disulfide linkages using N-succinimidyl-3-(2-
20 pyridylthio)propionate (SPDP) and succinimidyl 4-(N-maleimido-methyl)cyclohexane-1-carboxylate (SMCC) obtained from Pierce Company, Rockford, Illinois, (if the peptide lacks a sulfhydryl group, this can be provided by addition of a cysteine residue.) These reagents create a disulfide linkage between themselves and peptide cysteine residues on one protein and an
25 amide linkage through the epsilon-amino on a lysine, or other free amino group in the other. A variety of such disulfide/amide-forming agents are known. See, for example, Immun Rev (1982) 62:185. Other bifunctional coupling agents form a thioether rather than a disulfide linkage. Many of these thio-ether-forming agents are commercially available and include reactive esters of 6-maleimidocaproic acid,
30 2-bromoacetic acid, 2-iodoacetic acid, 4-(N-maleimido-methyl)cyclohexane-1-carboxylic acid, and the like. The carboxyl groups can be activated by combining them with succinimide or 1-hydroxyl-2-nitro-4-sulfonic acid, sodium salt. Additional methods of coupling antigens

WO 93/00365

-23-

PCT/US92/05388

employs the rotavirus/"binding peptide" system described in EPO Pub. No. 259,149, the disclosure of which is incorporated herein by reference. The foregoing list is not meant to be exhaustive, and modifications of the named compounds can clearly be used.

5 Any carrier may be used which does not itself induce the production of antibodies harmful to the host. Suitable carriers are typically large, slowly metabolized macromolecules such as proteins; polysaccharides, such as latex functionalized Sepharose®, agarose, cellulose, cellulose beads and the like; polymeric amino acids, such as polyglutamic acid, polylysine, and the like; amino
10 acid copolymers; and inactive virus particles, see, for example, Section II.D. Especially useful protein substrates are serum albumins, keyhole limpet hemocyanin, immunoglobulin molecules, thyroglobulin, ovalbumin, tetanus toxoid, and other proteins well known to those skilled in the art.

15 In addition to full-length viral proteins, polypeptides comprising

II.D. Preparation of Hybrid Particle Immunogens Containing HCV Epitopes

The immunogenicity of the epitopes of HCV may also be enhanced by preparing them in mammalian or yeast systems fused with or assembled with particle-forming proteins such as, for example, that associated with hepatitis B
20 surface antigen. See, e.g., US 4,722,840. Constructs wherein the HCV epitope is linked directly to the particle-forming protein coding sequences produce hybrids which are immunogenic with respect to the HCV epitope. In addition, all of the vectors prepared include epitopes specific to HBV, having various degrees of immunogenicity, such as, for example, the pre-S peptide. Thus, particles
25 constructed from particle forming protein which include HCV sequences are immunogenic with respect to HCV and HBV.

Hepatitis surface antigen (HBSAg) has been shown to be formed and assembled into particles in *S. cerevisiae* (P. Valenzuela *et al.* (1982)), as well as in, for example, mammalian cells (P. Valenzuela *et al.* (1984)). The formation of
30 such particles has been shown to enhance the immunogenicity of the monomer subunit. The constructs may also include the immunodominant epitope of HBSAg, comprising the 55 amino acids of the presurface (pre-S) region. Neurath *et al.* (1984). Constructs of the pre-S-HBSAg particle expressible in yeast are disclosed

WO 93/00365

-24-

PCT/US92/05388

in EPO 174,444, published March 19, 1986; hybrids including heterologous viral sequences for yeast expression are disclosed in EPO 175,261, published March 26, 1966. These constructs may also be expressed in mammalian cells such as Chinese hamster ovary (CHO) cells using an SV40-dihydrofolate reductase vector
5 (Michelle *et al.* (1984)).

In addition, portions of the particle-forming protein coding sequence may be replaced with codons encoding an HCV epitope. In this replacement, regions which are not required to mediate the aggregation of the units to form immunogenic particles in yeast or mammals can be deleted, thus eliminating
10 additional HBV antigenic sites from competition with the HCV epitope.

II.E. Preparation of Vaccines

Vaccines may be prepared from one or more immunogenic polypeptides derived from HCV. These polypeptides may be expressed in
15 various host cells (*e.g.*, bacteria, yeast, insect, or mammalian cells), or alternatively may be isolated from viral preparations or made synthetically. Single- or multi-valent vaccines against HCV may be comprised of one or more epitopes from one or more structural proteins, and/or one or more epitopes from one or more nonstructural proteins. These vaccines may be comprised of, for
20 example, recombinant HCV polypeptides and/or polypeptides isolated from the virions. In particular, vaccines are contemplated comprising one or more of the following HCV proteins, or subunit antigens derived therefrom: E1, E2, C, NS2, NS3, NS4 and NS5. Particularly preferred are vaccines comprising E1 and/or E2, or subunits thereof.

25 In addition to the above, it is also possible to prepare live vaccines of attenuated microorganisms which express one or more recombinant HCV polypeptides. Suitable attenuated microorganisms are known in the art and include, for example, viruses (*e.g.*, vaccinia virus (see Brown *et al.* (1986)), as well as bacteria.

30 The preparation of vaccines which contain an immunogenic polypeptide(s) as active ingredients, is known to one skilled in the art. Typically, such vaccines are prepared as injectables, either as liquid solutions or suspensions; solid forms suitable for solution in, or suspension in, liquid prior to injection may

WO 93/00365

-25-

PCT/US92/05388

also be prepared. The preparation may also be emulsified, or the protein encapsulated in liposomes. The active immunogenic ingredients are often mixed with excipients which are pharmaceutically acceptable and compatible with the active ingredient. Suitable excipients are, for example, water, saline, dextrose, glycerol, ethanol, or the like and combinations thereof. In addition, if desired, the vaccine may contain minor amounts of auxiliary substances such as wetting or emulsifying agents, pH buffering agents, and/or adjuvants which enhance the effectiveness of the vaccine. Examples of adjuvants which may be effective include but are not limited to: aluminum hydroxide, N-acetyl-muramyl-L-threonyl-D-isoglutamine (thr-MDP), N-acetyl-nor-muramyl-L-alanyl-D-isoglutamine (CGP 11637, referred to as nor-MDP), N-acetylmuramyl-L-alanyl-D-isoglutaminyl-L-alanine-2-(1'-2'-dipalmitoyl-sn-glycero-3-hydroxyphosphoryloxy)-ethylamine (CGP 19835A, referred to as MTP-PE), and RIBI, which contains three components extracted from bacteria, monophosphoryl lipid A, trehalose dimycolate and cell wall skeleton (MPL+TDM+CWS) in a 2% squalene/Tween® 80 emulsion. The effectiveness of an adjuvant may be determined by measuring the amount of antibodies directed against an immunogenic polypeptide containing an HCV antigenic sequence resulting from administration of this polypeptide in vaccines which are also comprised of the various adjuvants.

The vaccines are conventionally administered parenterally, by injection, for example, either subcutaneously or intramuscularly. Additional formulations which are suitable for other modes of administration include suppositories and, in some cases, oral formulations. For suppositories, traditional binders and carriers may include, for example, polyalkylene glycols or triglycerides; such suppositories may be formed from mixtures containing the active ingredient in the range of 0.5% to 10%, preferably 1%-2%. Oral formulations include such normally employed excipients as, for example, pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharine, cellulose, magnesium carbonate, and the like. These compositions take the form of solutions, suspensions, tablets, pills, capsules, sustained release formulations or powders and contain 10%-95% of active ingredient, preferably 25%-70%.

The proteins may be formulated into the vaccine as neutral or salt forms. Pharmaceutically acceptable salts include the acid addition salts (formed

WO 93/00365

-26-

PCT/US92/05388

with free amino groups of the peptide) and which are formed with inorganic acids such as, for example, hydrochloric or phosphoric acids, or such organic acids such as acetic, oxalic, tartaric, maleic, and the like. Salts formed with the free carboxyl groups may also be derived from inorganic bases such as, for example, sodium, potassium, ammonium, calcium, or ferric hydroxides, and such organic bases as isopropylamine, trimethylamine, 2-ethylamino ethanol, histidine, procaine, and the like.

II.F. Dosage and Administration of Vaccines

10 The vaccines are administered in a manner compatible with the dosage formulation, and in such amount as will be prophylactically and/or therapeutically effective. The quantity to be administered, which is generally in the range of 5 μ g to 250 μ g of antigen per dose, depends on the subject to be treated, capacity of the subject's immune system to synthesize antibodies, and the degree of protection desired. Precise amounts of active ingredient required to be administered may depend on the judgment of the practitioner and may be peculiar to each subject.

The vaccine may be given in a single dose schedule, or preferably in a multiple dose schedule. A multiple dose schedule is one in which a primary course of vaccination may be with 1-10 separate doses, followed by other doses given at subsequent time intervals required to maintain and or reenforce the immune response, for example, at 1-4 months for a second dose, and if needed, a subsequent dose(s) after several months. The dosage regimen will also, at least in part, be determined by the need of the individual and be dependent upon the judgment of the practitioner.

In addition, the vaccine containing the immunogenic HCV antigen(s) may be administered in conjunction with other immunoregulatory agents, for example, immune globulins.

30 II.G. Preparation of Antibodies Against HCV Epitopes

The immunogenic polypeptides described herein are used to produce antibodies, including polyclonal and monoclonal. If polyclonal antibodies are desired, a selected mammal (*e.g.*, mouse, rabbit, goat, horse, *etc.*) is immunized

WO 93/00365

-27-

PCT/US92/05388

with an immunogenic polypeptide bearing an HCV epitope(s). Serum from the immunized animal is collected and treated according to known procedures. If serum containing polyclonal antibodies to an HCV epitope contains antibodies to other antigens, the polyclonal antibodies can be purified by immunoaffinity chromatography. Techniques for producing and processing polyclonal antisera are known in the art, see for example, Mayer and Walker (1987). Alternatively, polyclonal antibodies may be isolated from a mammal which has been previously infected with HCV.

Monoclonal antibodies directed against HCV epitopes can also be readily produced by one skilled in the art. The general methodology for making monoclonal antibodies by hybridomas is well known. Immortal antibody-producing cell lines can be created by cell fusion, and also by other techniques such as direct transformation of B lymphocytes with oncogenic DNA, or transfection with Epstein-Barr virus. See, *e.g.*, M. Schreier *et al.* (1980); Hammerling *et al.* (1981); Kennett *et al.* (1980); see also, U.S. Patent Nos. 4,341,761; 4,399,121; 4,427,783; 4,444,887; 4,466,917; 4,472,500; 4,491,632; and 4,493,890. Panels of monoclonal antibodies produced against HCV epitopes can be screened for various properties; *i.e.*, for isotype, epitope affinity, *etc.*

Antibodies, both monoclonal and polyclonal, which are directed against HCV epitopes are particularly useful in diagnosis, and those which are neutralizing are useful in passive immunotherapy. Monoclonal antibodies, in particular, may be used to raise anti-idiotypic antibodies.

Anti-idiotypic antibodies are immunoglobulins which carry an "internal image" of the antigen of the infectious agent against which protection is desired. See, for example, Nisonoff, A., *et al.* (1981) and Dreesman *et al.* (1985). Techniques for raising anti-idiotypic antibodies are known in the art. See, for example, Grzych (1985), MacNamara *et al.* (1984), and Vytdehaag *et al.* (1985). These anti-idiotypic antibodies may also be useful for treatment, vaccination and/or diagnosis of NANBH, as well as for an elucidation of the immunogenic regions of HCV antigens.

II.H. Immunoassay and Diagnostic Kits

Both the polypeptides and the antibodies of the present invention are useful in immunoassays to detect presence of HCV antibodies, or the presence of

WO 93/00365

-28-

PCT/US92/05388

the virus and/or HCV polypeptides (or epitopes), in, for example, biological samples. Design of the immunoassays is subject to a great deal of variation, and many formats are known in the art. The immunoassay will utilize at least one viral epitope derived from HCV. In one embodiment, the immunoassay uses a combination of viral epitopes derived from HCV. These epitopes may be derived from the same or from different viral polypeptides, and may be in separate recombinant or natural polypeptides, or together in the same recombinant polypeptides. An immunoassay may use, for example, a monoclonal antibody directed towards a viral epitope(s), a combination of monoclonal antibodies directed towards epitopes of one viral antigen, monoclonal antibodies directed towards epitopes of different viral antigens, polyclonal antibodies directed towards the same viral antigen, or polyclonal antibodies directed towards different viral antigens. Protocols may be based, for example, upon competition, or direct reaction, or sandwich type assays. Protocols may also, for example, use solid supports, or may be by immunoprecipitation. Most assays involve the use of labeled antibody or polypeptide; the labels may be, for example, enzymatic, fluorescent, chemiluminescent, radioactive, or dye molecules. Assays which amplify the signals from the probe are also known; examples of which are assays which utilize biotin and avidin, and enzyme-labeled and mediated immunoassays, such as ELISA assays (described below).

Typically, an immunoassay for an anti-HCV antibody(s) will involve selecting and preparing the test sample suspected of containing the antibodies, such as a biological sample, then incubating it with an antigenic (*i.e.*, epitope-containing) HCV polypeptide(s) under conditions that allow antigen-antibody complexes to form, and then detecting the formation of such complexes. Suitable incubation conditions are well known in the art. The immunoassay may be, without limitations, in a heterogenous or in a homogeneous format, and of a standard or competitive type.

In a heterogeneous format, the polypeptide is typically bound to a solid support to facilitate separation of the sample from the polypeptide after incubation. Examples of solid supports that can be used are nitrocellulose (*e.g.*, in membrane or microtiter well form), polyvinyl chloride (*e.g.*, in sheets or microtiter wells), polystyrene latex (*e.g.*, in beads or microtiter plates,

WO 93/00365

-29-

PCT/US92/05388

polyvinylidene fluoride (known as Immulon®), diazotized paper, nylon membranes, activated beads, and Protein A beads. For example, Dynatech Immulon® 1 or Immulon® 2 microtiter plates or 0.25 inch polystyrene beads (Precision Plastic Ball) can be used in the heterogeneous format. The solid support containing the antigenic polypeptide is typically washed after separating it from the test sample, and prior to detection of bound antibodies. Both standard and competitive formats are known in the art.

In a homogeneous format, the test sample is incubated with antigen in solution. For example, it may be under conditions that will precipitate any antigen-antibody complexes which are formed. Both standard and competitive formats for these assays are known in the art.

In a standard format, the amount of HCV antibodies forming the antibody-antigen complex is directly monitored. This may be accomplished by determining whether labeled anti-xenogenic (*e.g.*, anti-human) antibodies which recognize an epitope on anti-HCV antibodies will bind due to complex formation. In a competitive format, the amount of HCV antibodies in the sample is deduced by monitoring the competitive effect on the binding of a known amount of labeled antibody (or other competing ligand) in the complex.

Complexes formed comprising anti-HCV antibody (or, in the case of competitive assays, the amount of competing antibody) are detected by any of a number of known techniques, depending on the format. For example, unlabeled HCV antibodies in the complex may be detected using a conjugate of antixenogeneic Ig complexed with a label, (*e.g.*, an enzyme label).

In immunoassays where HCV polypeptides are the analyte, the test sample, typically a biological sample, is incubated with anti-HCV antibodies under conditions that allow the formation of antigen-antibody complexes. Various formats can be employed. For example, a "sandwich assay" may be employed, where antibody bound to a solid support is incubated with the test sample; washed; incubated with a second, labeled antibody to the analyte, and the support is washed again. Analyte is detected by determining if the second antibody is bound to the support. In a competitive format, which can be either heterogeneous or homogeneous, a test sample is usually incubated with antibody and a labeled,

WO 93/00365

-30-

PCT/US92/05388

competing antigen is also incubated, either sequentially or simultaneously. These and other formats are well known in the art.

Efficient detection systems for HCV infection may include the use of panels of epitopes, as described above. The epitopes in the panel may be
5 constructed into one or multiple polypeptides. The assays for the varying epitopes may be sequential or simultaneous.

The enzyme-linked immunosorbent assay (ELISA) can be used to measure either antigen or antibody concentrations. This method depends upon conjugation of an enzyme to either an antigen or an antibody, and uses the bound
10 enzyme activity as a quantitative label. To measure antibody, the known antigen is fixed to a solid phase (*e.g.*, a microplate or plastic cup), incubated with test serum dilutions, washed, incubated with anti-immunoglobulin labeled with an enzyme, and washed again. Enzymes suitable for labeling are known in the art, and include, for example, horseradish peroxidase. Enzyme activity bound to the
15 solid phase is measured by adding the specific substrate, and determining product formation or substrate utilization colorimetrically. The enzyme activity bound is a direct function of the amount of antibody bound.

To measure antigen, a known specific antibody is fixed to the solid phase, the test material containing antigen is added, after an incubation the solid
20 phase is washed, and a second enzyme-labeled antibody is added. After washing, substrate is added, and enzyme activity is estimated colorimetrically, and related to antigen concentration. Kits suitable for immunodiagnosis and containing the appropriate labeled reagents are constructed by packaging the appropriate materials, including the polypeptides of the invention containing HCV epitopes or
25 antibodies directed against HCV epitopes in suitable containers, along with the remaining reagents and materials required for the conduct of the assay, as well as a suitable set of assay instructions.

III. General Methods

30 The general techniques used in the practice of the present invention can be found in, for example, the references cited herein, particularly EPO Pub. Nos. 318,216 and 388,232, as well as the references in the bibliography, which are incorporated herein by reference.

WO 93/00365

-31-

PCT/US92/05388

IV. Examples

Described below are examples of the present invention which are provided only for illustrative purposes, and not to limit the scope of the present invention. In light of the present disclosure, numerous embodiments within the scope of the claims will be apparent to those of ordinary skill in the art.

IV.A. Epitope Mapping of HCV Genome

The following example is the result of an epitope mapping experiment conducted on the HCV1 polyprotein sequence shown in Fig. 1. As shown in Fig. 3-11, there are heterogeneities among HCV isolates, indicating that these amino acid substitutions can be made in the octamers described below. In addition to substitutions with amino acids from the corresponding location in other HCV isolates, substitutions with synthetic analogs of the particular amino acids or conservative substitutions based on charge, etc. (particularly when the substitution does not destroy antibody binding) are within the scope of the invention.

IV.A.1. Synthesis of Overlapping Peptides

Polyethylene pins arranged on a block in an 8×12 array (Coselco Mimetopes, Victoria, Australia) were prepared by placing the pins in a bath (20% v/v piperidine in dimethylformamide (DMF)) for 30 minutes at room temperature. The pins were then removed, washed in DMF for 5 min, then washed in methanol four times (2 min/wash). The pins were allowed to air dry for at least 10 min, then washed a final time in DMF (5 min). 1-Hydroxybenzotriazole (HOBt, 367 mg) was dissolved in DMF (80 mL) for use in coupling Fmoc-protected amino acids: Fmoc-L-Ala-OPfp, Fmoc-L-Cys(Trt)-OPfp, Fmoc-L-Asp(O-*t*Bu)-OPfp, Fmoc-L-Glu(O-*t*Bu)-OPfp, Fmoc-L-Phe-OPfp, Fmoc-Gly-OPfp, Fmoc-L-His(Boc)-OPfp, Fmoc-L-Ile-OPfp, Fmoc-L-Lys(Boc)-OPfp, Fmoc-L-Leu-OPfp, Fmoc-L-Met-OPfp, Fmoc-L-Asn-OPfp, Fmoc-L-Pro-OPfp, Fmoc-L-Gln-OPfp, Fmoc-L-Arg(Mtr)-OPfp, Fmoc-L-Ser(*t*-Bu)-ODhbt, Fmoc-L-Thr(*t*-Bu)-ODhbt, Fmoc-L-Val-OPfp, and Fmoc-L-Tyr-OPfp.

The protected amino acids were placed in microtiter plate wells with HOBt, and the pin block placed over the plate, immersing the pins in the wells. The assembly was then sealed in a plastic bag and allowed to react at 25°C for 18

WO 93/00365

-32-

PCT/US92/05388

hours to couple the first amino acids to the pins. The block was then removed, and the pins washed with DMF (2 min), MeOH (4×2 min), and again with DMF (2 min) to clean and deprotect the bound amino acids. The procedure was repeated for each additional amino acid coupled, until all octamers had been prepared.

The free N-termini were then acetylated to compensate for the free amide, as most of the epitopes are not found at the N-terminus and thus would not have the associated positive charge. Acetylation was accomplished by filling the wells of a microtiter plate with DMF/acetic anhydride/triethylamine (5:2:1 v/v/v) and allowing the pins to react in the wells for 90 min at 20°C. The pins were then washed with DMF (2 min) and MeOH (4×2 min), and air dried for at least 10 min.

The side chain protecting groups were removed by treating the pins with trifluoroacetic acid/phenol/dithioethane (95:2.5:2.5, v/v/v) in polypropylene bags for 4 hours at room temperature. The pins were then washed in dichloromethane (2×2 min), 5% di-isopropylethylamine/dichloromethane (2×5 min), dichloromethane (5 min), and air-dried for at least 10 min. The pins were then washed in water (2 min), MeOH (18 hours), dried *in vacuo*, and stored in sealed plastic bags over silica gel.

20

IV.A.2. Assay of Peptides

Octamer-bearing pins prepared as above were first treated by sonicating for 30 min in a disruption buffer (1% sodium dodecylsulfate, 0.1% 2-mercaptoethanol, 0.1 M NaH_2PO_4) at 60°C. The pins were then immersed several times in water (60°C), followed by boiling MeOH (2 min), and allowed to air dry. The pins were then precoated for 1 hour at 25°C in microtiter wells containing 200 μL blocking buffer (1% ovalbumin, 1% BSA, 0.1% Tween®, and 0.05% NaN_3 in PBS), with agitation. The pins were then immersed in microtiter wells containing 175 μL antisera obtained from human patients diagnosed as having HCV and allowed to incubate at 4°C overnight. The pins were assayed against antisera from three individual patients. Specimen #PAA 3663-s ("A") exhibited strong reaction to HCV Western blots, HCV competitive ELISA, HCV ELISA to clone C100-3 (at 1:1000 dilution), and RIBA responses of >4+ to C100, 5-1-1,

30

WO 93/00365

-33-

PCT/US92/05388

and C33c (C22 not done). (Antigen/clone names are per EPO Pub. Nos. 318,216 and 388,232, as well as those described in the literature regarding HCV immunoassays available from Ortho Diagnostics Systems, Inc.) Neat plasma was diluted 1:500 in blocking buffer. Specimen #PAA 33028 ("B") exhibited strong
5 reaction to HCV Western blots, HCV competitive ELISA, HCV ELISA to clone C100-3 (at 1:500 dilution), and RIBA responses of >4+ to C100, 5-1-1, C33C and C22. Polyclonal antisera was partially purified by passage through a protein A column, and was used at a dilution of 1:200 in blocking buffer. Specimen #PAA s32931 ("C") exhibited moderate reaction to HCV Western blots (3+),
10 HCV competitive ELISA, HCV ELISA to clone C100-3 (at 1:64 dilution), and RIBA responses of 3+ and 4+ to C100 and 5-1-1, respectively (C33c and C22 not done). Polyclonal antisera was partially purified by passage through a protein A column, and was used at a dilution of 1:500 in blocking buffer.

The pins were washed in PBS/Tween® 20 (4 × 10 min) at room
15 temperature, then incubated in microtiter wells containing horseradish peroxidase-labeled goat anti-Human Ig antisera (175 µL, 1:2000 dilution in blocking buffer without NaN₃) for 1 hour at 25°C with agitation. The antihuman antisera is specific for human Ig light and heavy chains, and reacts with both IgG and IgM classes. The pins were again washed in PBS/Tween® 20 (4 × 10 min) at room
20 temperature. Substrate solution was prepared by diluting NaH₂PO₄ (1 M, 200 mL) and citric acid (1 M, 160 mL) to 2 L with distilled water, adjusting the pH to 4.0. Azino-di-3-ethylbenzthiazodinsulfonate (ABTS, 50 mg) and hydrogen peroxide (0.3 µL/mL) was added to 100 mL of buffer immediately prior to use to complete the substrate solution. The substrate solution (150 µL) was added to each well of
25 a microtiter plate, and the pins immersed in the wells and incubated at 25°C in the dark. After color developed, the reactions were halted by removing the pins, and absorbance of the solutions read at 405 nm.

The octamers listed below were immunoreactive with anti-HCV antisera. Peptides reacting with all three antisera are listed as epitopes, while peptides reacting with only one or two antisera are listed as weak epitopes (indicated
30 by "~"). Particularly strong epitopes are labeled with letters rather than numbers (e.g., EpAA).

WO 93/00365

-34-

PCT/US92/05388

	<u>AA#</u>	<u>Sequence</u>		<u>AA#</u>	<u>Sequence</u>
5	23	KEPGGGQA	Ep2	87	GNEGCGWA
	24	EPGGGQAV		88	NEGCGWAG
	25	PGGGQAVG		89	EGCGWAGW
	26	GGGQAVGG		90	GCGWAGWL
	27	GGQAVGGV		91	CGWAGWLL
	28	GQAVGGVY		92	GWAGWLLS
10	29	QAVGGVYL ~		93	WAGWLLSP
	30	AVGGVYLL		94	AGWLLSPR
	31	VGGVYLLP		95	GWLLSPRG
	32	GGVYLLPR		96	WLLSPRGS
	33	GVYLLPRR		97	LLSPRGSR
	34	VYLLPRRG		98	LSPRGSRP
15	35	YLLPRRGP		99	SPRGSRPS
	36	LLPRRGPR			
20	66	PKARRPEG Ep1	100	PRGSRPSW Ep3	
	67	KARRPEGR	101	RGSRPSWG	
	68	ARRPEGRT	102	GSRPSWGP	
	69	RRPEGRTW	103	SRPSWGPT	
	70	RPEGRTWA			
	71	PEGRTWAQ	186	TVPASAYQ Ep4	
25	72	EGRTWAQP	187	VPASAYQV	
	73	GRTWAQPG EpA	188	PASAYQVR	
	74	RTWAQPGY	189	ASAYQVRN	
	75	TWAQPGYP	190	SAYQVRNS	
	76	WAQPGYPW	191	AYQVRNST	
	77	AQPGYPWP			
30	78	QPGYPWPL	206	DCPNSSIV ~	
	79	PGYPWPLG			
	80	GYPWPLYG	223	TPGCVPCV ~	
	81	YPWPLYGN			
	82	PWPLYGNE	232	EGNASRCW Ep5	
	83	WPLYGNEG			
35	84	PLYGNEGC	256	TQLRRHID ~	
	85	LYGNEGCG			
	86	YGNEGCGW	286	LVGQLFTF ~	
40			297	RHWTQGC Ep6	
			298	HWTQGCN	
			299	WTTQGCNC	
45			321	MMMNWSPI ~	
			347	DMLAGAHW Ep7	
			357	LAGIAYFS Ep8	

WO 93/00365

-35-

PCT/US92/05388

	413	LINTNGSW EpB	594	YSRCGSGP F _A
	414	INTNGSWH	595	SRCGSGPW
			596	RCGSGPWL
	432	SLNTGWLA ~	597	CGSGPWLT
5			598	GSGPWLT
	465	FDQGWGPI EpC	599	SGPWLT
	466	DQGWGPIS	600	GPWLTPRC
	467	QGWGPISY	601	PWLTPRCL
	468	GWGPISYA	602	WLTPRCLV
10	469	WGPISYAN	603	LTPRCLVD EpF
	470	GPISYANG	604	TPRCLVDY
	471	PISYANGS	605	PRCLVDYP
			606	RCLVDYPY
	480	PDQRPYCW EpD	607	CLVDYPYR
15	481	DQRPYCW	608	LVDYPYRL
	482	QRPYCW	609	VDYPYRLW
	483	RPYCW	610	DYPYRLWH
	484	PYCWHYP	611	YPYRLWHY F _B
			612	PYRLWHYP
20	500	KSVCGPVY Ep9	613	YRLWHYPC
	501	SVCGPVYC		
	502	VCGPVYCE	641	EAACNWTR
	521	RSGAPTYS Ep10		
25			662	LSPLLLII Ep13
	540	NNTRPPLG EpE	663	SPLLLII
	541	NTRPPLGN	664	PLLLIIQ
	542	TRPPLGNW	665	LLLLIIQW
	543	RPPLGNWF		
30	544	PPLGNWFG	685	LSTGLIHL ~
	545	PLGNWFGC		
	546	LGNWFGCT	705	VGSSIASW ~
	547	GNWFGCTW	706	GSSIASWA
	548	NWFGCTWM		
35	549	WFGCTWMN	729	ARVCSCIW Ep14
	579	LHCPTDCF ~	782	WVPGAVYT
40			783	VPGAVYTF
			784	PGAVYTFY
			785	GAVYTFYG
			786	AVYTFYGM
			787	VYTFYGMW
			788	YTFYGMWP
			789	TFYGMWPL
45				
			801	LALPQRAY ~

Ep12

EpG

WO 93/00365

-36-

PCT/US92/05388

5	851	RVEAQLHV EpH	1384	VIKGRHL Ep20
	852	VEAQLHVW	1410	LGINAVAY Ep21
	853	EAQLHVWI	1411	GINAVAYY
	854	AQLHVWIP	1454	CNTCVIQT ~
	855	QLHVWIPP	1492	GRGKPGIY Ep22
10	893	LAVFGPLW ~	1493	RGKPGIYR
	916	QGLLRFCA ~	1532	PAETTVRL Ep23
	928	MIGGHYVQ Ep15	1533	AETTVRLR
	946	TGTYVYNH EpI	1534	ETTVRLRA
			1535	TTVRLRAY
15	952	NHLTPLRD EpJ	1560	GVFIGLIH Ep24
	953	HLTPLRDW	1561	VFIGLIHI
	954	LTPLRDWA	1581	ENLPYLVA
20	1026	LAPITAYA ~	1567	NLPYLVAY
	1072	TCINGVCW EpK	1568	LPYLVAYQ
	1109	LVGWPAPO Ep16	1569	PYLVAYQA
25	1171	CPAGHAVG Ep17	1570	YLVAYQAT
	1113	PAGHAVGI	1571	LVAYQATV
	1114	AGHAVGIF	1572	VAYQATVC
	1115	GHAVGIFR	1573	AYQATVCA
	1116	HAVGIFRA	1574	YQATVCAR
	1117	AVGIFRAA	1575	QATVCARQ
30	1218	VVPQSEQV EpL	1576	ATVCARQA
	1240	VPAAYAAQ ~	1577	TVCARQAP
	1260	ATLGFGAY ~	1601	PPSWDQMW
35	1280	GVRITTGS Ep18	1602	PSWDQMWK
	1281	VRITTGSP	1603	SWDQMWKC
	1282	RITTGSP	1604	WDQMWKCL
40	1283	ITTGSPIT	1605	DQMWKCLI
	1284	TTGSPITY	1606	QMWKCLIR
	1285	TGSPITYG	1607	MWKCLIRL
	1322	DATSILGI ~	1615	KPTLHGPI Ep27
	1338	TAGARLVV ~	1616	PTLHGPIP
45			1617	TLHGPIPL
			1618	LHGPIPLL
			1619	HGPIPLLY
	1371	GEIPFYGK Ep19	1620	GPIPLLYR
			1655	VVTSTWVL ~

WO 93/00365		-37-	PCT/US92/05388	
5	1694	IIPDREVL EpM	1966	SECTIPCS EpS
	1695	IPDREVL	1967	ECTIPCSG
	1710	ECSQHLPY EpN	1968	CTIPCSGS
	1711	CSQHLPYI	1969	TIPCSGSW
	1712	SQHLPYIE		
10	1728	EKQKALGL EpO	1999	LMPQLPGI EpT
	1729	KQKALGLL	2000	MPQLPGIP
	1758	EIEWAKLM EpP	2001	PQLPGIPE
	1759	IEWAKLMW	2002	QLPGIPEV
	1760	EWAKLMWN	2003	LPGIPEVS
15	1761	WAKLMWNE	2004	PGIPEVSC
	1762	AKLMWNEI	2005	GIPEVSCQ
	1781	LPGNPAIA ~	2006	IPEVSCQR
	1808	LFNILGGW Ep28	2007	PEVSCQRG
	1821	AAPGAATA ~	2008	EVSCQRGY
20	1851	ILAGYGAG Ep29	2009	VSCQRGYK
	1880	VNLLPAIL ~	2010	SCQRGYKG
	1908	PGEGAVQW EpG	2011	CQRGYKGV
	1909	GEGAVQWM	2012	QRGYKGVW
	1910	EGAVQWMN	2013	RGYKGVWR
30	1911	GAVQWMNR	2014	GYKGVWRG
	1912	AVQWMNRL		
	1913	VQWMNRLI		
	1925	RGNHVSPI EpR	2024	IMHTRCHC
	1940	AAARVTAI Ep30		
35	1941	AARVTAI	2048	VGPRICRN EpU
	1942	ARVTAIS	2049	GPRICRNY
	1943	RVTAILSS	2050	PRICRNYW
	1944	VTAILSSL	2051	RICRNYWS
	1945	TAILSSLV	2052	ICRNYWSG
40	1946	AILSSLVT	2053	CRNYWSGT
	1947	ILSSLVTQ	2054	RNYWSGTE
	1948	LSSLVTQL	2055	NYWSGTEP
	1949	SSLVTQLL	2056	YWSGTEPI
	1950	SLVTQLLR	2057	WSGTEPIN
45	1951	LVTQLLR	2071	TPLPAPNY Ep32
			2088	EEYVIRQV EpV
			2089	EYVIRQVG
			2090	YVIRQVGD
			2091	VIRQVGDF
			2092	IRQVGDFH
			2093	RQVGDFHY
			2108	DNLKCPCQ ~

Ep31

WO 93/00365

-38-

PCT/US92/05388

5	2122	IELDGVVR EpW	2280	REAQALPV EpZ
	2123	IELDGVRL	2281	EAQALPVW
	2124	ELDGVRLH	2282	AQALPVWA
	2125	LDGVRLHR	2283	QALPVWAR
	2126	DGVRLHRF	2284	ALPVWARP
10	2127	GVRLHRFA	2285	LPVWARPD
	2128	VRLHRFAP	2286	PVWARPDY
	2129	RLHRFAPP	2287	VWARPDYN
	2130	LHRFAPPC	2288	WARPDYNP
	2131	HRFAPPCK	2289	ARPDYNPP
15	2132	RFAPPCKP	2290	RPDYNPPL
	2133	FAPPCKPL		
	2134	APPCKPLL	2325	PPPRKKRT Ep35
	2135	PPCKPLL	2326	PPRKKRTV
	2136	PCKPLLRE	2327	PRKKRTVV
20	2137	CKPLLREE		
	2138	KPLLREEV	2345	AELASRSE Ep36
	2139	PLLREEVS	2346	ELASRSEG
	2140	LLREEVSF EpX	2347	LASRSEGS
	2141	LREEVSFR	2348	ASRSEGSS
25	2142	REEVSFRV	2349	SRSEGSSS
	2143	EEVSFRVG		
	2144	EVSFRVGL	2382	AESYSSMP Ep37
	2145	VSFRVGLH		
	2146	SFRVGLHE	2401	SDGSWSTV
30	2147	FRVGLHEY	Ep38	
	2148	RVGLHEYP		
	2165	EPEPDVAV ~	EpAA	
	2187	GRRLARGS -		
	2226	LIBANLLW EpY	2417	VVCCSMSY
35	2227	IEANLLWR	2418	VCCSMSYW
	2228	EANLLWRQ	2419	CCSMSYWI
	2229	ANLLWRQE	2420	CSMSYWIG
	2230	NLLWRQEM	2421	SMSYWIGA
	2231	LLWRQEMG	2422	MSYWIGAL
40	2232	LWRQEMGG		
	2244	VESENKVV Ep33		
	2245	ESENKVVI		
	2246	SENKVVIL		
	2247	ENKVVILD		
45	2248	NKVVILDS		
	2249	KVVILDSF		
	2250	VVILDSFD		
	2267	EISVPAEI Ep34		

WO 93/00365

-39-

PCT/US92/05388

5	2439	QKLPINAL EpBB	EpDD	2602	LPLAVMGS
	2440	KLPINALS		2603	PLAVMGSS
	2441	LPINALSN		2604	LAVMGSSY
	2442	PINALSNS		2605	AVMGSSYG
	2443	INALSNSL		2606	VMGSSYGE
10	2444	NALSNSLL	Ep41	2607	MGSSYGEQ
	2445	ALSNSLLR		2608	GSSYGEQR
	2446	LSNSLLRH		2609	SSYGEQRV
	2447	SNSLLRHH		2610	SYGEQRVE
	2448	NSLLRHHN		2611	YGEQRVEE
15	2449	SLLRHHNL	Ep41	2612	GEQRVEEL
	2450	LLRHHNLV		2632	KTPMGFSY
	2451	LRHHNLVY		2633	TPMGFSYD
	2452	RHHNLVYS		2634	PMGFSYDT
	2453	HHNLVYST		2635	MGFSYDTR
20	2454	HNLVYSTI	EpEE	2636	GFSYDTRC
	2455	NLVYSTIS		2637	FSYDTRCE
	2456	LVYSTISR		2638	SYDTRCED
	2469	QKKVTFDR Ep39		2660	YQCCDLDP ~
	2470	KKVTFDRL		2676	LTERLYVG
25	2471	KVTFDRLQ	EpEE	2677	TERLYVGG
	2472	VTFDRLQV		2678	ERLYVGGP
	2473	TFDRLQVL		2679	RLYVGGPL
	2474	FDRLQVLD		2688	NSRGENCEG
	2475	DRLQVLDS		2689	SRGENCGY
30	2476	RLQVLDSH	EpFF	2690	RGENCGYR
	2495	ASKVKANL ~		2691	GENCGYRR
	2533	RKAVTHIN EpCC		2692	ENCGYRRC
	2534	KAVTHINS		2693	NCGYRRCR
	2573	GRKPARLI Ep40		2707	TSCGNTLI Ep42
35	2574	RKPARLIV	Ep43	2721	AACRAAGL ~
	2575	KPARLIVF		2757	AFTEAMTR
	2576	PARLIVFP		2758	FTEAMTRY
	2577	ARLIVFPD		2759	TEAMTRYSA
	2578	RLIVFPDL		2760	EAMTRYSA
40			Ep43	2761	AMTRYSA
				2762	MTRYSA
45			Ep43		

WO 93/00365

-40-

PCT/US92/05388

5	Ep45	2779	DLELIISC Ep44	2878	DLPPIIQR Ep47
		2780	LELIISC	2879	LPPIIQRL
		2794	HDGAGKRV	2880	PPIIQR LH
		2795	DGAGKRVY	2881	PIIQR LHG
		2796	GAGKRVYY	2882	IIQR LHGL
10		2797	AGKRVYYL	2883	IQR LHGLS
		2798	GKRVYYLT	2884	QRLHGLSA
		2799	KRVYYLTR	2885	RLHGLSAF
		2800	RVYYLTRD	2886	LHGLSAFS
		2801	VYYLTRDP	2887	HGLSAFSL
15	EpGG	2802	YYLTRDPT	2888	GLSAFSLH
		2817	WETARHTP	2889	LSAFSLHS
		2818	ETARHTPV	2890	SAFSLHSY
		2819	TARHTPVN	2891	AFSLHSYS
		2820	ARHTPVNS	2892	FSLHSYSP
20		2821	RHTPVNSW	2893	SLHSYSPG
		2822	HTPVNSWL	2894	LHSYSPGE
		2823	TPVNSWLG	2895	HSYSPGEI
		2824	PVNSWLGNI		
		2825	VNSWLGNI		
25		2826	NSWLGNI		
		2827	SWLGNIIM		
		2828	WLGNIIME		
		2829	LGNIMEA		
		2830	GNIMEAP		
30		2831	NIIMEAPT		
		2832	IIMEAPTL		
		2833	IMEAPTLW		
		2834	MEAPTLWA		
		2835	EAPTLWAR		
35		2836	APTLWARM		
		2837	PTLWARM		
		2838	TLWARMIL		
		2839	LWARMILM		
		2840	WARMILMT		
40		2841	ARMILMTH		
		2842	RMILMTHF		
		2843	MILMTHFE		
45		2863	LDCEIYGA Ep46		
		2864	DCEIYGAC		
		2865	CEIYGACY		
		2866	EIYGACYS		
		2867	IYGACYSI		

WO 93/00365

-41-

PCT/US92/05388

	2912	LGVPPLRA Ep48
	2913	GVPPLRAW
	2914	VPPLRAWR
5	2938	AAICGKYL Ep49
	2939	AICGKYLE
	2940	ICGKYLEN
10	2966	DLSGWETA
	2984	VSHARPRW EpII

15 IV.B. Differential Assay

The following assay was performed to distinguish early antigens from later antigens. Antibodies to the early antigens may be detected, and used to diagnose HCV infection more quickly.

Serial bleeds were obtained from a human patient presenting with elevated ALT, but negative for anti-C100-3 antibody. The five bleeds obtained prior to complete seroconversion (C100-3 positive) were pooled and used in the assay at a dilution of 1:2000. The assay was conducted as described in Section IV.A. above. However, one duplicate set of pins was incubated with horseradish peroxidase-labeled goat anti-Human IgG specific antisera, while the other set was incubated with horseradish peroxidase-labeled goat anti-Human IgM specific anti-
 25 sera. Epitopes immunoreactive with IgM antibodies are early epitopes.

The results indicated that most early epitopes are found in the region extending from about amino acid 480 to about amino acid 650. Particularly strong IgM epitopes were octamers beginning with amino acid nos. 506, 510, 523, 553, 562, 580, and the region from 590 to 620. Assays which employ antigens bearing epitopes from this region will permit diagnosis of HCV infection at an
 30 early point than assays employing other antigens.

We have additionally tested serial plasma specimens taken from five patients with open heart post-transfusion NANB hepatitis, with studies followed for 3-12 years. Initial bleed dates were less than one week apart. Each specimen was tested for IgG and IgM by EIA against one core antigen (C22), two envelope antigens (E1 and E2), and three nonstructural region antigens (C33c, C100, and
 35

WO 93/00365

-41-

PCT/US92/05388

2912 LGVPPLRA Ep48
 2913 GVPPLRAW
 2914 VPPLRAWR

 5 2938 AAICGKYL Ep49
 2939 AICGKYLE
 2940 ICGKYLEN

 10 EpHH 2966 DLSGWETA

 2984 VSHARPRW EpII

15 IV.B. Differential Assay

The following assay was performed to distinguish early antigens from later antigens. Antibodies to the early antigens may be detected, and used to diagnose HCV infection more quickly.

20 Serial bleeds were obtained from a human patient presenting with
 elevated ALT, but negative for anti-C100-3 antibody. The five bleeds obtained
 prior to complete seroconversion (C100-3 positive) were pooled and used in the
 assay at a dilution of 1:2000. The assay was conducted as described in Section
 IV.A. above. However, one duplicate set of pins was incubated with horseradish
 peroxidase-labeled goat anti-Human IgG specific antisera, while the other set was
 25 incubated with horseradish peroxidase-labeled goat anti-Human IgM specific anti-
 sera. Epitopes immunoreactive with IgM antibodies are early epitopes.

The results indicated that most early epitopes are found in the
 region extending from about amino acid 480 to about amino acid 650. Particularly
 strong IgM epitopes were octamers beginning with amino acid nos. 506, 510, 523,
 30 553, 562, 580, and the region from 590 to 620. Assays which employ antigens
 bearing epitopes from this region will permit diagnosis of HCV infection at an
 early point than assays employing other antigens.

We have additionally tested serial plasma specimens taken from five
 patients with open heart post-transfusion NANB hepatitis, with studies followed
 35 for 3-12 years. Initial bleed dates were less than one week apart. Each specimen
 was tested for IgG and IgM by EIA against one core antigen (C22), two envelope
 antigens (E1 and E2), and three nonstructural region antigens (C33c, C100, and

WO 93/00365

-42-

PCT/US92/05388

NS5). We found that the IgM response to C22 and C33c preceeded the IgG response for those antigens. NS-5 also induced an IgM response, but this response did not preceed the IgG response for that antigen. Thus, one can prepare assays capable of determining very early stages of infection by utilizing epitopes
5 derived from the C22 and C33c regions and assaying for IgM binding. Antibodies to the C33c region persisted for the longest period of time, suggesting that diagnostic assays directed toward C33c should be the most reliable.

IV.C. Sequence Variations in HCV Isolates from Different Individuals

10 Isolates of HCV which contain sequences which deviate from CDC/HCV1 were identified in human individuals, some of whom were serologically positive for anti-C100-3 antibodies (EC10 was antibody negative). Identification of these new isolates was accomplished by cloning and sequencing segments of the HCV genome which had been amplified by the PCR technique
15 using CDC/HCI sequences. The method utilizes primers and probes based upon the HCV cDNA sequences described herein. The first step in the method is the synthesis of a cDNA to either the HCV genome, or its replicative intermediate, using reverse transcriptase. After synthesis of the HCV cDNA, and prior to amplification, the RNA in the sample is degraded by techniques known in the art.
20 A designated segment of the HCV cDNA is then amplified by the use of the appropriate primers. The amplified sequences are cloned, and clones containing the amplified sequences are detected by a probe which is complementary to a sequence lying between the primers, but which does not overlap the primers.

25 IV.C.1. HCV Isolates Isolated from Humans in the U.S.

Blood samples which were used as a source of HCV virions were obtained from the American Red Cross in Charlotte, North Carolina, and from the Community Blood Center of Kansas, Kansas City, Missouri. The samples were screened for antibodies to the HCV C100-3 antigen using an ELISA assay and
30 subjected to supplemental Western blot analysis using a polyclonal goat anti-human HRP to measure anti-HCV antibodies. Two samples, #23 and #27, from the American Red Cross and from the Community Blood Center of Kansas, respectively, were determined to be HCV positive by these assays.

WO 93/00365

-43-

PCT/US92/05388

Viral particles present in the serum of these samples were isolated by ultracentrifugation under the conditions described by Bradley *et al.* (1985). RNA was extracted from the particles by digestion with proteinase K and SDS at final concentrations of 10 $\mu\text{g/mL}$ proteinase K, and 0.1% SDS; digestion was for
5 1 hour at 37°C. Viral RNA was further purified by extraction with chloroform-phenol.

HCV RNA in the preparation of RNA was reverse transcribed into cDNA. After both strands of the cDNA were synthesized, the resulting cDNA was then amplified by the PCR method. The HCV cDNAs in three clones derived
10 from each HCV isolate, were subjected to sequence analysis. Analysis was essentially by the method described in Chen and Seeburg (1985).

Consensus sequences of the clones derived from HCV in samples 23 and 27 are shown in Fig. 3 and Fig. 4, respectively. The variable sequences are also shown in these figures, as are the amino acids encoded in the consensus
15 sequences.

Figures 5 and 6 show comparisons of the aligned positive strand nucleotide sequences (Fig. 5) and putative amino acid sequences (Fig. 6) of samples 23, 27, and HCV1. The amino acid sequence of HCV1 in Fig. 6 represents amino acid numbers 129-467 of the HCV polyprotein encoded by the
20 large ORF in the HCV genomic RNA. An examination of Figs. 5 and 6 show that there are variations in the sequences of the three isolated clones. The sequence variations at the nucleotide level and the amino acid level are summarized in the table immediately below. In the table, the polypeptides designated S and NS1 represent amino acid numbers 130 to ~380, and 380 to ~470, respectively, as
25 those domains were previously known. The numbering is from the putative initiator methionine. The terminology S and NS1 is based upon the positioning of the sequences encoding the polypeptides using the Flavivirus model. As discussed above, however, recent evidence suggests that there is not total correlation between HCV and the Flaviviruses with regard to viral polypeptide domains,
30 particularly in the putative E/NS1 domains. Indeed, HCV polypeptides and their coding domains may exhibit substantial deviation from the Flavivirus model.

WO 93/00365

-44-

PCT/US92/05388

TABLE:

		<u>Sequence Homology</u>					
		Nucleotide Encoding			Amino Acid Encoded		
		overall	S	NS1	overall	S	NS1
5		%	%	%	%	%	%
	HCV1/HCV23	93	95	91	92	95	87
	HCV1/HCV27	89	93	84	89	95	82
	HCV23/HCV27	89	93	85	90	93	84

10 Although there are variations in the newly isolated HCV sequences, the cloned sequences from samples 23 and 27 (called HCV23 and HCV27) each contain 1019 nucleotides, indicating a lack of deletion and addition mutants in this region in the selected clones. The sequences in Figs. 5 and 6 also show that the isolated sequences are not rearranged in this region.

15 A comparison of the consensus sequences for HCV1 and for the other isolates of HCV is summarized in the Table, *supra*. The sequence variations between the chimpanzee isolate HCV1, and the HCVs isolated from humans are about the same as that seen between the HCVs of human origin.

20 It is of interest that the sequence variations in two of the putative domains is not uniform. The sequence in a putative S region appears to be relatively constant, and randomly scattered throughout the region. In contrast, a putative NS1 region has a higher degree of variability than the overall sequence, and the variation appears to be in a hypervariable pocket of about 28 amino acids which is located about 70 amino acids downstream from the putative N-terminus
25 of the putative polyprotein.

 Although it may be argued that the detected variations were introduced during the amplification process, it is unlikely that all of the variations are from this result. It has been estimated that Taq polymerase introduces errors into a sequence at approximately one base per 10 kilobases of DNA template per
30 cycle (Saiki *et al.* (1988)). Based upon this estimate, up to 7 errors may have been introduced during the PCR amplification of the 1019 bp DNA fragment. However, the three subclones of HCV-23 and HCV-27 yielded 29 and 14 base variations, respectively. The following suggest that these variations are naturally

WO 93/00365

-45-

PCT/US92/05388

occurring. About 60% of the base changes are silent mutations which do not change the amino acid sequence. Variations introduced by the Taq polymerase during PCR amplification would be expected to occur randomly; however, the results show that the variant sequences are clustered in at least one specific region.

5

IV.C.2. HCV Isolates from Humans in Italy and in the U.S.

Segments of HCV RNA present in different isolates were amplified by the HCV/cPCR method. These segments span a region of ~0.6 Kb to ~1.6 Kb downstream from the methionine encoding start codon of the putative HCV polyprotein. The isolates are from biological specimens obtained from HCV infected individuals. More specifically, isolate HCT #18 is from human plasma from an individual in the U.S.A., EC1 and EC10 are from a liver biopsy of an Italian patient, and Th is from a peripheral blood mononucleocyte fraction of an American patient. Comparable segments of HCV RNA have been isolated from a chimpanzee.

15

RNA was extracted from the human plasma specimens using phenol:CHCl₃:isoamyl alcohol extraction. Either 0.1 mL or 0.01 mL of plasma was diluted to a final volume of 1.0 mL, with a TENB/proteinase K/SDS solution (0.05 M Tris-HCL, pH 8.0, 0.001 M EDTA, 0.1 M NaCl, 1 mg/mL Proteinase K, and 0.5 % SDS) containing 10 to 40 µg/mL polyadenylic acid, and incubated at 37°C for 60 minutes. After this proteinase K digestion, the resultant plasma fractions were deproteinized by extraction with TE (50 mM Tris-HCl, pH 8.0, 1 mM EDTA) saturated phenol, pH 6.5. The phenol phase was separated by centrifugation, and was reextracted with TENB containing 0.1 % SDS. The resulting aqueous phases from each extraction were pooled, and extracted twice with an equal volume of phenol/chloroform/isoamyl alcohol [1:1(99:1)], and then twice with an equal volume of a 99:1 mixture of chloroform/isoamyl alcohol. Following phase separation by centrifugation, the aqueous phase was brought to a final concentration of 0.2 M Na acetate, and the nucleic acids were precipitated by the addition of two volumes of ethanol. The precipitated nucleic acids were recovered by ultracentrifugation in a SW 41 rotor at 38 K, for 60 minutes at 4°C or in a microfuge for 10 minutes at 10 K, 4°C.

25

30

WO 93/00365

-46-

PCT/US92/05388

RNA extracted from the liver biopsy was provided by Dr. F. Bonino, Ospedale Maggiore di S. Giovanni Battista, Torino, Italy. The mononucleocyte fraction was obtained by sedimentation of the individual's aliquot of blood through Ficoll-Paque® (Pharmacia Corp), using the manufacturer's directions. Total RNA was extracted from the fraction using the guanidinium thiocyanate procedure described in Choo et al (1989).

Synthesis of HCV cDNA from the samples was accomplished using reverse transcriptase. Following ethanol precipitation, the precipitated RNA or nucleic acid fraction was dried, and resuspended in DEPC treated distilled water. Secondary structures in the nucleic acids were disrupted by heating at 65°C for 10 minutes, and the samples were immediately cooled on ice. cDNA was synthesized using 1 to 3 µg of total RNA from liver, or from nucleic acids (or RNA) extracted from 10 to 100 µL of plasma. The synthesis utilized reverse transcriptase, and was in a 25 µL reaction, using the protocol specified by the manufacturer, BRL. All reaction mixtures for cDNA synthesis contained 23 units of the RNAase inhibitor, Rnasin® (Fisher/Promega). Following cDNA synthesis, the reaction mixtures were diluted with water, boiled for 10 minutes, and quickly chilled on ice.

Each set of samples was subjected to two rounds of PCR amplification. The primers for the reactions were selected to amplify regions designated "EnvL" and EnvR". The "EnvL" region encompasses nucleotides 669-1243, and putative amino acids 117 to 308; the "EnvR" region encompasses nucleotides 1215-1629, and encodes putative amino acids 300-408 (the putative amino acids are numbered starting from the putative methionine initiation codon).

The PCR reactions were performed essentially according to the manufacturer's directions (Cetus-Perkin-Elmer), except for the addition of 1 µg of RNase A. The reactions were carried out in a final volume of 100 µL. The PCR was performed for 30 cycles, utilizing a regimen of 94°C (1 min), 37°C (2 min), and 72°C (3 min), with a 7 minute extension at 72°C for the last cycle. The samples were then extracted with phenol:CHCl₃, ethanol precipitated two times, resuspended in 10 mM Tris HCl, pH 8.0, and concentrated using Centricon-30 (Amicon) filtration. This procedure efficiently removes oligonucleotides less than

WO 93/00365

-47-

PCT/US92/05388

30 nucleotides in size; thus, the primers from the first round of PCR amplification are removed.

The Centricon-30 concentrated samples were then subjected to a second round of PCR amplification. Amplification by PCR was for 35 cycles
5 utilizing a regimen of 94°C (1 min), 60°C (1 min), and 72°C (2 min), with a 7 minute extension at 72°C for the last cycle. The samples were then extracted with phenol:CHCl₃, precipitated two times, and digested with EcoRI. The PCR reaction products were analyzed by separation of the products by electrophoresis on 6% polyacrylamide gels. DNA of approximately the estimated size of the
10 expected PCR product was electroeluted from the gels, and subcloned into either a pGEM-4 plasmid vector or into λgt11. The expected product sizes for the EnvL and EnvR after the first round of amplification are 615 bp and 683 bp, respectively; after the second round of amplification the expected product sizes for EnvL and EnvR are 414 bp and 575 bp, respectively. The plasmids containing the
15 amplified products were used to transform host cells; the pGEM-4 plasmid was used to transform DH5-alpha, and λgt11 was used to transform C600 delta-HFL. Clones of the transformed cells which either hybridized to the appropriate HCV probes, or those which had inserts of the correct size were selected. The inserts were then cloned in M13 and sequenced. The probes for all of the HCV/cPCR
20 products consisted of ³²P labeled sections of HCV cDNA which had been prepared by PCR amplification.

Sequence information on variants in the EnvL region was obtained from 3 clones from HCT #18, 2 clones from TH, 3 clones from EC1, and from the HCV1 clones. A comparison of the composite nucleotide sequence of each
25 isolate derived from these clones is shown in Fig. 7. In the figure, each sequence is shown 5' to 3' for the sense strand for the EnvL region, and the sequences have been aligned. The vertical lines and capital letters indicate sequence homology, the absence of a line and an uncapitalized letter indicates a lack of homology. The sequences shown in the lines are as follows: line 1, Thorn; line 2, EC1; line 3,
30 HCT #18; line 4, HCV1.

Sequence information on variants in the EnvR region was obtained from two clones of EC10, and from HCV1 clones. The two EC10 clones differed by only one nucleotide. A comparison of the nucleotide sequences of EC10(clone

WO 93/00365

-48-

PCT/US92/05388

2) and a composite of the HCV1 sequences is shown in Fig. 8; each sequence is shown 5' to 3' for the sense strand of the EnvR region, and the sequences have been aligned. The double dots between the sequences indicate sequence homology.

- 5 A comparison of the amino acid sequences encoded in the EnvL (amino acids #117-308) and EnvR region (amino acids #300-438) for each of the isolates is shown in Fig. 9 and Fig. 10, respectively. Included in the Figures are sequences for the isolates JH23 and JH27, described above. Also indicated are sequences from a Japanese isolate; these sequences were provided by Dr. T. Miyamura, Japan. In the figures, the amino acid sequence for the region is given in its entirety for HCV1, and the non-homologous amino acids in the various isolates are indicated.

As seen in Fig. 9, In the EnvL region there is overall about a 93% homology between HCV1 and the other isolates. HCT18, Th, and EC1 have about a 97% homology with HCV1; JH23 and JH27 have about 96% and about 95% homology, respectively, with HCV1. Fig. 10 shows that the homologies in the EnvR region are significantly less than in the EnvL region; moreover, one subregion appears to be hypervariable (*i.e.*, from amino acid 383-405). This data is summarized in the Table immediately below.

20 Table: Homology of EnvR Region

Isolate	Percent Homology with HCV1	
	AA330-AA438	AA383-AA405
JH23(U.S.)	83	57
JH27(U.S.)	80	39
25 Japanese	73	48
EC10 (Italy)	84	48

VI. Industrial Applicability

- The epitopes identified herein can be used to make polypeptide products as described above for applications such as the screening of blood for HCV infection, clinical HCV diagnosis, the generation of antibodies, and preparation of medicaments. Other applications are described above, and still others will be readily apparent to those of ordinary skill.
- 30

VII. Bibliography

- Barr *et al.*, Biotechniques (1986) 4:428.
- Bradley *et al.*, Gastroenterology (1985) 88:773.
- Botstein, Gene (1979) 8:17.
- 5 M.A. Brinton, (1986) in "The Viruses: The Togaviridae and Flaviviridae" (Series eds. Fraenkel-Conrat and Wagner, vol. eds. Schlesinger and Schlesinger, Plenum Press), p.327-374.
- Broach (1981) in: "Molecular Biology of the Yeast - *Saccharomyces*, Vol. 1, p.445, Cold Spring Harbor Press.
- 10 Broach *et al.*, Meth Enzymol (1983) 101:307.
- Catty (1988), "Antibodies, Volume 1: A Practical Approach" (IRL Press).
- Chaney *et al.*, Cell Mol Genet (1986) 12:237.
- Chakrabarti *et al.*, Mol Cell Biol (1985) 5:3403.
- Chang *et al.*, Nature (1977) 198:1056.
- 15 Chen and Seeburg, DNA (1985) 4:165.
- Chirgwin *et al.*, Biochemistry (1979) 18:5294.
- Chomczynski and Sacchi, Anal Biochem (1987) 162:156.
- Choo *et al.*, Science (1989) 244:359.
- Clewell *et al.*, Proc Natl Acad Sci USA (1969) 62:1159.
- 20 Clewell, J Bacteriol (1972) 110:667.
- Cohen, Proc Natl Acad Sci USA (1972) 69:2110.
- Cousens *et al.*, Gene (1987) 61:265.
- De Boer *et al.*, Proc Natl Acad Sci USA (1983) 292:128.
- Dreesman *et al.*, J Infect Dis (1985) 151:761.
- 25 S.M. Feinstone and J.H. Hoofnagle, New Engl J Med (1984) 311:185.
- Felgner *et al.*, Proc Natl Acad Sci USA (1987) 84:7413.
- Fields & Knipe (1986), "Fundamental Virology" (Raven Press, N.Y.).
- Fiers *et al.*, Nature (1978) 273:113.
- R.J. Gerety *et al.*, in "Viral Hepatitis and Liver Disease" (B.N. Vyas, J.L.
- 30 Dienstag, and J.H. Hoofnagle, eds)
- Glennie *et al.*, Nature (1982) 295:712.
- Gluzman, Cell (1981) 23:175.
- Goeddel *et al.*, Nuc Acids Res (1980) 8:4057.

WO 93/00365

-50-

PCT/US92/05388

- Graham and Van der Eb, Virology (1978) 52:546.
- Grunstein and Hogness, Proc Natl Acad Sci USA (1975) 73:3961.
- Grych *et al.*, Nature (1985) 316:74.
- Gubler and Hoffman Gene (1983) 25:263.
- 5 Hahn, Virology (1988) 162:167.
- Hammerling *et al.*, (1981), "Monoclonal Antibodies and T-Cell Hybridomas".
- Han, Biochemistry (1987) 26:1617.
- Helfman, Proc Natl Acad Sci USA (1983) 80:31.
- Hess *et al.*, J Adv Enzyme Reg (1968) 7:149.
- 10 Hinnen *et al.*, Proc Natl Acad Sci USA (1978) 75:1929.
- Hitzeman *et al.*, J Biol Chem (1980) 255:2073.
- Holland *et al.*, Biochemistry (1978) 17:4900.
- Holland, J Biol Chem (1981) 256:1385.
- Holland and Holland, J Biol Chem (1980) 255:2596.
- 15 Hoopes *et al.*, Nuc Acids Res (1981) 9:5493.
- Houghton *et al.*, Nuc Acids Res (1981) 9:247
- T.V. Hunyh *et al.*, (1985) in "DNA Cloning Techniques; A Practical Approach" (D. Glover, Ed., IRL Press, Oxford, U.K.) pp. 49-78.
- Immun Rev (1982) 62:185.
- 20 Ito *et al.*, Agric Biol Chem (1984) 48:341.
- Iwarson, British Medical J (1987) 295:946.
- Kennett *et al.* (1980) "Monoclonal Antibodies".
- Kniskern *et al.*, Gene (1986) 46:135.
- Kyte and Doolittle, J Mol Bio (1982) 157:105-132.
- 25 Laemmli, Nature (1970) 227:680.
- Lee *et al.*, Science (1988) 239:1288.
- Luckow and Summers, Virol (1989) 17:31.
- Mackett *et al.*, J Virol (1984) 49:857.
- T. Maniatis *et al.* (1982) "Molecular Cloning; A Laboratory Manual" (Cold
- 30 Spring Harbor Press, Cold Spring Harbor, N.Y.).
- Sambrook *et al.* (1989), "Molecular Cloning; A Laboratory Manual", Second Edition (Cold Spring Harbor Press, Cold Spring Harbor, N.Y.).
- Manning and Mocarski, Virol (1988) 167:477.

WO 93/00365

-51-

PCT/US92/05388

- Mayer and Walker, eds. (1987), "Immunochemical Methods In Cell And Molecular Biology" (Academic Press, London).
- Maxam *et al.*, Meth Enzymol (1980) 65:499.
- MacNamara *et al.*, Science (1984) 226:1325.
- 5 Messing *et al.*, Nuc Acids Res (1981) 9:309.
- Messing, Meth Enzymol (1983) 101:20-37.
- Michelle *et al.*, Int. Symposium on Viral Hepatitis.
- Monath (1986) in "The Viruses: The Togaviridae And Flaviviridae" (Series eds. Fraenkel-Conrat and Wagner, vol. eds. Schlesinger and Schlesinger, Plenum
- 10 Press), pp. 375-440.
- Moss (1987) in "Gene Transfer Vectors For Mammalian Cells" (Miller and Calos, eds., Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.), p. 10.
- Nagahuma *et al.*, Anal Biochem (1984) 141:74.
- Neurath *et al.*, Science (1984) 224:392.
- 15 Nisonoff *et al.*, Clin Immunol Immunopathol (1981) 21:397-406.
- L.R. Overby, Curr Hepatol (1985) 5:49.
- L.R. Overby, Curr Hepatol (1986) 6:65.
- L.R. Overby, Curr Hepatol (1987) 7:35.
- Pachl *et al.*, J Virol (1987) 61:315.
- 20 Peleg, Nature (1969) 221:193.
- Pfefferkorn and Shapiro (1974), in "Comprehensive Virology", Vol. 2 (Fraenkel-Conrat & Wagner, eds., Plenum, N.Y.) pp. 171-230.
- A.M. Prince, Ann Rev Microbiol (1983) 37:217.
- Rice *et al.*, Science (1985) 229:726.
- 25 Rice *et al.* (1986) in "The Viruses: The Togaviridae And Flaviviridae" (Series eds. Fraenkel-Conrat and Wagner, vol. eds. Schlesinger and Schlesinger, Plenum Press), p.279-328.
- Roehrig (1986) in "The Viruses: The Togaviridae And Flaviviridae" (Series eds. Fraenkel-Conrat and Wagner, vol. eds. Schlesinger and Schlesinger, Plenum
- 30 Press)
- Sadler *et al.*, Gene (1980) 8:279.
- Saiki *et al.*, Nature (1986) 324:163.
- Saiki *et al.*, Science (1988) 239:487.

WO 93/00365

-52-

PCT/US92/05388

- Sanger *et al.*, Proc Natl Acad Sci USA (1977) 74:5463.
- Setlow, ed. (1988), "Genetic Engineering" Vol. 10, pp. 195-219 (Plenum Publishing Co., N.Y.).
- Schlesinger *et al.*, J Virol (1986) 60:1153.
- 5 M. Schreier *et al.*, (1980) "Hybridoma Techniques"
- Scopes (1984), "Protein Purification, Principles and Practice", 2nd Ed., (Springer-Verlag, N.Y.).
- Shimatake *et al.*, Nature (1981) 292:128.
- Singh *et al.*, Nuc Acids Res (1983) 11:4049.
- 10 Sippel, Eur J Biochem (1973) 37:31.
- Smith *et al.*, Mol Cell Biol (1983) 3:2156-2165.
- Steimer *et al.*, J Virol (1986) 58:9.
- Stollar (1980), in "The Togaviruses" (R.W. Schlesinger, ed.), pp. 584-622.
- Stuve *et al.*, J Virol (1987) 61:326.
- 15 Sumiyoshi *et al.*, Virology (1987) 161:497.
- Taylor *et al.*, Biochim Biophys Acta (1976) 442:324.
- Towbin *et al.*, Proc Natl Acad Sci USA (1979) 76:4350.
- Tsu and Herzenberg (1980), in "Selected Methods In Cellular Immunology" (W.H. Freeman and Co.) pp. 373-391.
- 20 Vytdehaag *et al.*, J Immunol (1985) 134:1225.
- P. Valenzuela *et al.*, Nature (1982) 298:344.
- P. Valenzuela *et al.*, (1984), in "Hepatitis B" (I. Millman *et al.*, ed, Plenum Press) pp. 225-236.
- Warner, DNA (1984) 3:401.
- 25 Ward *et al.*, Nature (1989) 341:544.
- Wu and Grossman, Meth Enzymol (1987), 154 "Recombinant DNA", Part E.
- Wu, Meth Enzymol (1987), 155, "Recombinant DNA", part F.
- Zoller, Nuc Acids Res (1982) 10:6487.
- U.S. Patent No. 4,341,761 U.S. Patent No. 4,466,917
- 30 U.S. Patent No. 4,399,121 U.S. Patent No. 4,472,500
- U.S. Patent No. 4,427,783 U.S. Patent No. 4,491,632
- U.S. Patent No. 4,444,887 U.S. Patent No. 4,493,890
- U.S. Patent No. 4,816,467

WO 93/00365

-53-

PCT/US92/05388

WHAT IS CLAIMED:

1. A polypeptide comprising a truncated HCV sequence containing an HCV epitope of the formula

5

 aa_x-aa_y

wherein aa denotes an amino acid;

x and y are integers such that $y-x \geq 6$;

aa_x-aa_y indicates a portion of the amino acid sequence of Figure 1; and

10 x is selected from the group consisting of 23-34, 36, 66-79, 81-94, 96-98, 101-103, 186-189, 191, 206, 223, 232, 256, 286, 297-299, 321, 347, 357, 413, 414, 432, 465-471, 480-484, 501, 502, 521, 540-549, 579, 594-599, 601-613, 641, 662-665, 685, 705, 706, 729, 782-789, 801, 851-855, 893, 916, 928, 946, 952-954, 1026, 1072, 1109, 1112-1117, 1218, 1240, 1280-1285, 1322, 1338, 1371, 1384, 1410, 1411, 1454, 1492, 1493, 1532-1535, 1560, 1561, 1566-1568, 1571-1577, 1601-1607, 1615-1620, 1655, 1695, 1710-1712, 1728, 1729, 1758-1762, 1781, 1808, 1821, 1851, 1880, 1908-1913, 1925, 1940-1948, 1951, 1966-1969, 1999, 2001-2004, 2006-2014, 2024, 2048-2053, 2055-2057, 2071, 2088-2093, 2108, 2122-2148, 2165, 2187, 2226-2232, 2244-2249, 2267, 2281-2286, 2288, 2289, 2325-2327, 2346, 2347, 2349, 2382, 2401, 2417-2422, 2439-2444, 20 2446-2456, 2469, 2471-2476, 2495, 2533, 2534, 2573-2578, 2602-2604, 2606-2612, 2632-2638, 2660, 2676-2679, 2688-2693, 2707, 2721, 2757-2762, 2779, 2794, 2795, 2797-2799, 2801, 2802, 2817-2843, 2863-2867, 2878-2884, 2886-2895.

25

2. The polypeptide of claim 1 which is about 100 amino acids or less in length.

3. The polypeptide of claim 2 wherein $y-x \leq 50$.

30

4. The polypeptide of claim 2 wherein $y-x \leq 20$.

5. The polypeptide of claim 2 wherein $y-x < 10$.

WO 93/00365

-54-

PCT/US92/05388

6. The polypeptide of claim 2 wherein x is selected from the group consisting of 506, 510, 523, 553, 562, 580, and 590-620.

7. A polypeptide of about 100 amino acids or less comprising an
5 HCV epitope of the formula



wherein aa denotes an amino acid;

x and y are integers such that $y-x \geq 6$;

aa_x-aa_y indicates a portion of the amino acid sequence of Figure 1; and

10 x is selected from the group consisting of 35 (where y is less than 45),
80 (where y is less than 90), 95 (where y is less than 110), 99 (where y is less
than 120), 100 (where y is less than 150), 190 (where y is less than 210), 500
(where y is less than 550), 600 (where y is less than 625), 1260 (where y is less
than 1280), 1569 (where y is less than 1931), 1570 (where y is less than 1590),
15 1694 (where y is less than 1735), 1949 (where y is less than 2124), 1950 (where y
is less than 1985), 2000 (where y is less than 2050), 2005 (where y is less than
2025), 2054 (where y is less than 2223), 2250 (where y is less than 2330), 2287
(where y is less than 2385), 2290 (where y is less than 2310), 2345 (where y is
less than 2375), 2348 (where y is less than 2464), 2445 (where y is less than
20 2475), 2470 (where y is less than 2490), 2605 (where y is less than 2620), 2780
(where y is less than 2830), 2796 (where y is less than 2886), 2800 (where y is
less than 2850), and 2885 (where y is less than 2905).

8. An immunoassay reagent comprising a polypeptide according to
25 any of claims 1-7.

9. The immunoassay reagent of claim 8 wherein $y-x \leq 50$.

10. The immunoassay reagent of claim 8 wherein x is selected from
30 the group consisting of 506, 510, 523, 553, 562, 580, and 590-620.

WO 93/00365

-55-

PCT/US92/05388

11. A method for detecting the presence of antibodies immunoreactive with Hepatitis C virus (HCV) proteins in a sample, said method comprising:
contacting an immobilized immunoassay reagent according to claim 8 with said sample; and
5 detecting antibodies bound to said reagent.

12. A method for inducing an immunological response in a subject against HCV, said method comprising:
10 administering to said subject an effective amount of a polypeptide according to any one of claims 1-7.

13. A composition for inducing an immunological response in a subject against HCV, said composition comprising an effective amount of a
15 polypeptide according to any one of claims 1-7.

14. A monoclonal or polyclonal antibody composition wherein said antibodies bind the HCV epitope of a polypeptide according to claim 1.

20 15. A method of making a polypeptide according to any of claims 1-7 wherein said polypeptide is prepared by recombinant expression or chemical synthesis.

WO 93/00365

PCT/US92/05388

1 / 23

FIG. 1 - 1

R T

MSTNPKPQKKNRNTNRRPQDVKFPGGGOIVGGVYLLPRRGPRLGVRATR
KTSERSQPRGRROPIPKARRPEGRTWAQPGYPWPLYGNEGCGWAGWLLSP-100
RGSRP SWGPTDPRRRSRNLGKVIDTLTCGFADLMGYIPLVGAPLGGAARA

T

LAHGV RVLEDGVNYATGNLPGCSFSIFLLALLSCLTVPASAYQVRNSTGL-200
YHVTNDCPNSSIVYEADAILHTPGCVPCVREGNASRCWVAMTPTVATRD
GKLPA TQLRRHIDLLVGSATLCSALYVGDLGSGVFLVGQLFTFSPRRHWT-300

V

TQGCNCSIYPGHITGHRMAWDMMNWSPTTALVMAQLLRIPQAILDMIAG
AHWGVLAGIAYFSMVGNWAKVLVLLLLFAGVDAETHVTGGSAGHTVSGFV-400
SLLAPGAKQNVQLINTNGSWHLNSTALNCNDSLNTGWLGLFYHHKFNSS
GCPERLASCRPLTDFDQGWGPISYANGSGPDQRPYCWHYPPKPCGIVPAK-500
SVC GPVYCFTPSPVVVGTTDRSGAPTYSWGENDTDVFLNNTRPPLGNWF
GCTWMNSTGFTKVC GAPPCVIGGAGNNTLHCPTDCFRKHPPDATYSRCGSG-600

I

PWLTFRCLVDYPYRLWHYPCTINYTIFKIRMYVGGVEHRLEAACNWSTRGE
RCDLED RDRSELSPLLLTTTQWQVLPCSFTTLPALSTGLIHLHQNIQVDVQ-700
YLYGVGSSIASWAIKWEYVLLFLLLADARVCSCLWMMLLISQAEAALEN
LVILNAASLAGTHGLVSFLVFFCFAYLKGKWVPGAVYTFYGMWPLLLLL-800

WO 93/00365

PCT/US92/05388

2 / 23

(N)
LALPORAYALDTEVAASC GG VVLVGLMAL TLSPYYKRYISWCLWWLOYFL
TRVEAQLHVWIPPLNVRGGRDAVILLMCAVHPTLVFDITKLLLAVFGPLW-900
ILQASLLKVPHYFVRVQGLLRFCALARKMIGGHYVQMVI IKLGALTGTIVY
NHLTPLRDWAHNGLRDLAVAVEPVVFSQMETKLITWGADTAACGDIINGL-1000
PVSARRGREILLGPADGMVSKGWRL LAPITAYAQQTRGLLGCIITSLTGR
DKNQVEGEVQIVSTAAQTFLATCINGVCWTVYHGAGTRTIASPKGPVIQM-1100

S T
YTNVDQDLVGWPAPQGSRSRLTPCTCGSSDLYLVTRHADVIPVRRRGDSRG
SLLSPRPISYLGSSGGPLLCPAGHAVGIFRAAVCTRGVAKAVDFIPVEN-1200
LETTMRSPVFTDNSSPPVVPQSFQVAHLHAPTGS GKSTKVPAAAYAAQGYK

L
VLVLNPSVAATLGFGAYMSKAHGIDPNIRTGVRTITTGSPITYSTYGKFL-1300

ADGGCSGGAYDIIICDECHSTDATSILGIGTVLDQAETAGARLVVLATAT
PPGSVTVPHPNIEEVALSTTGEIPFYGKAIPLEVIKGRHLIFCHSKKKK-1400
DELA AKLV ALGINAVAYYRGLDVSVIPTSGDVVVVATDALMTGYTGDFDS

Y (S)
VIDCNTCVTQTVD FSLDPTFTIETITLPQDAVSRTQRRGRTGRGKPGIYR-1500
FVAPGERPSGMFDSSVLCECYDAGCAWYELTPAETTVRLRAYMNTPLPV
CQDHLEFWEGVFTGLTHIDAHFLSQTQSGENLPYLVAYQATVCARAQAP-1600
PPSWDOMWKCLIRLKPTLHGPTPLLYRLGAVQNEITLTHPVTKYIMTMS
ADLEVVTSTWVLVGGVLAALAAAYCLSTGCVVIVGRVVLSGKPAIIPDREV-1700
LYREFDEMEEC SQHLPYIEQGMMLAEQFKQKALGLLQTASROAEVIAPAV
QTNWQKLETFWAKHMWNFISGIQYLAGLSTLPGNPAIASLMAFTA AVTSP-1800
LTTSQTL LFNI LGGWVAAQLAAPGAATAFVGAGLAGAAIGSVGLGKVLID

(G)
ILAGYGAGVAGALVAFKIMSGEVPSTEDLVNLLPAILSPGALVVGVC AA-1900

(HC)
ILRRHVGPGE GAVQWMNRLIAFASRGNHVSPTHYVPESDAAARVTAILSS

FIG. 1-2

WO 93/00365

PCT/US92/05388

3 / 23

LTVTQLLRRLHQWISSECTTPCSGSWLRDIWDWICEVLSDFKTWLKAKLM-2000

(V)

PQLPGIPFVSCQRGYKGVWRGDGIMHTRCHCGAEITGHVKNGTMRIVGPR
TCRNMWSGTFPINAYTTGPCTPLPAPNYTFALWRVSAEEYVEIROVGDFH-2100
YVTGMTTDNLKCPCQVPSPEFFTELDGVRLLHRFAPPCKPLLREEVSFRVG
LHEYFVGSQLPCEPEPDVAVLTSMLTDPSSHITAEAAAGRRLARGSPPSVAS-2200
SSASQLSAPSLKATCTANHDSFDAELIEANLLWRQEMGGNITRVESENKV
VILDSFDPLVAEEDEREISVPAEILRKSRRFAQALPVWARPDYNPPLVET-2300

(S)

WKKPDYEPVHVHGCPLPPPKSPPVPPPRKKRTTVVLTESTLSTALAEATR

(FA)

SFGSSSTSGITGDNTTTSSEPAPSGCPPDSDAESYSSMPPLEGEPGDPDL-2400
SDGSWSTVSSEANAEDVCCSMSYSWTGALVTPCAAEQKLPINALSNSL
LRHHNLVYSTTSRSACQRQKKVTFDRLQVLD SHYQDVLKEVKAASKVKA-2500

(F)

NLLSVEEACSLTPPHSAKSKFGYGAKDVRCHARKAVTHINSVWKDLLEDN
VTPIDTTIMAKNEVFCVQPEKGRKPARLIVFPDLGVRVCEKMALYDVVT-2600
KLPLAVMGSSYGFOYSPGQRYEFLVQAWKSKKTPMGFSYDTRCFDSTVTE

(G)

SDIRTEEAIYQCCDLDPQARVAIKSLTERLYVGGPLTNSRGENCYRRRCR-2700
ASGVLTTSCGNTLTICYIKARAACRAAGLQDCTMLVCGDDL VICESAGVQ
EDAASLRAFTEAMTRYSAAPPDPPQPEYDLELITSCSSNVSVAHGAGKR-2800
VYYLTRDPTTPLARAAWETARHTPVNSWLGNIIIMFAPTLWARMILMTHFF
SVLIARDQLEQALDCEIYGACYSIEPLDLPPIIQRLHGLSAFSLHSYSPG-2900

G

EINRVAACLRKLGVPPLRAWRHRARSVRARLLARGGRAAICGKYLFNWAV

(P)

RTKLKLTPIAAAGQLDLSGWFTAGYSGGDIYHSVSHARPRWIWFCLLLLA-3000
AGVGIIYLLPNRO-3011

Stop codon


() = Heterogeneity due possibly
to 5' or 3' terminal cloning
artefact.


FIG. 1-3

WO 93/00365

PCT/US92/05388

4 / 23





WO 93/00365

PCT/US92/05388

6 / 23

FIG. 2-3

AGCGCAACTGCACGTGTGGATTCCCCCTCAACGTCCGAGGGGGCGGACGCCGTGCAT
 CTTACTCATGTGTGTGTACACCCGACTCTGGTATTGACATCACCAAAATTGCTGTCTGGC-3000
 CGTCTTCGGACCCCTTTGGATTCTTCAAGCCAGTTTGCTTAAAGTACCCCTACTTTGTGCG
 CGTCCAAGGCCCTTCTCCGGTTCTGCGGTTAGCGGGAAGATGATCGGAGGCCATTACGT
 GCAAATGGTCAATCAATTAAGTTAGGGCGCTTACTGGCACCTATGTTTATAACCATCTCAC
 TCCTCTTCGGGACTGGGCGCACAAAGGCTTTCGAGATCTGGCCGTGGCTGTAGAGCCAGT
 CGTCTTCTCCCAATGAGAGACCAAGCTCATACGTGGGGGCGAGATACCGCCGCTGCGG-3300
 TGACATCATCAACGGCTTGCCCTGTTTCCGCCCGAGGGCGGGAGATACTGCTCGGGCC
 AGCCGATGGAATGGTCTCCAAAGGGTGGAGTTGCTGGCGCCCATCACGGCGTACGCCCA
 GCAGACAAGGGCCCTCTAGGGTGCAATAATCACAGCCTAACTGGCCGGGACAAACCA
 AGTGGAGGGTGAGGTCAGATTGTGTCAACTGCTGCCCAACCTTCTCTGGCAACGTGCAT
 CAATGGGGTGTGCTGACTGTCTATACCAGGGGCGGGAACGAGGACCATCGCGTCACCCAA-3600

T

GGGTCCCTGTCAATCCAGATGTATACCAATGTAGACCAAGACCTTGTGGGTGGCCCGCTCC

C

GCAAGGTAGCCGCTCATTTGACACCCCTGCACCTTGCGGCTCCTCGGACCTTTACCTGGTCAC
 GAGGCACGCCGATGTCAATCCCGTGCGCCGGGGGTGATAGCAGGGCAGCCTGCTGTC
 GCCCGGCCCATTTCTTACTTGAAGGCTCTCGGGGTCGCTGTTGTGCCCGCGGG
 GCACGCCGTGGCATAATTAGGGCCGCGGTGTGCACCCGTGGAGTGGCTAAGGCGGTGA-3900
 CTTTATCCCTGTGGAGAACCTAGAGACAACCATGAGTCCCCGGTGTTCACGGATAACTC
 CTCGCCACAGTAGTGCCCCAGAGCTTCCAGGTGGCTCACCTCCATGCTCCACAGGCAG
 CGGCAAAAGCACCAAGGTCCCGCTGCATATGCAGCTCAGGGCTATAAGGTGCTAGTACT
 CAACCCCTCTGTGTGCAACACTGGGCTTGGTGCTTACATGTCCAAGGCTCATGGGAT

T

CGATCCTAACATCAGGACCGGGGTGAGAACAAATTACCACTGGCAGCCCCATCAGTACTC-4200
 CACCTACGGCAAGTTCTTGCCGACGGCGGTGCTCGGGGGCGCTTATGACATAATAAT
 TTGTGACGAGTGCCACTCCACGGATGCCACATCCATCTTGGGCATCGGCACCTGCTTGA
 CCAAGCAGAGACTGCGGGGGGAGACTGGTTGTGCTCGCCACCGCCACCCCTCCGGGCTC
 CGTCACTGTGCCCATCCCAACATCGAGGAGTTGCTCTGTCCACCAACCGGAGATCCC
 TTTTACGGCAAGGCTATCCCCCTCGAAGTAATCAAGGGGGGAGACATCTCATCTCTG-4500
 TCATTCAAAGAAGTAGTCGACGAACCTCGCCGCAAGCTGGTCGCAATTGGGCATCAATGC
 CGTGGCCCTACTACCGCGGTCTTGACGTGTCCGTCATCCCGACCAAGCGGGATGTTGTCTG

WO 93/00365

PCT/US92/05388

7 / 23

FIG.2-4

A

CGTGGCAACCGATGCCCTCATGACCGGCTATACCGGGACTTCGACTCGGTGATAGACTG
 CAATACGTGTGTCACCCAGACAGTCCGATTTACGCCCTTGACCCCTACCTTCACCAATTGAGAC
 AATACGCTCCCCAGGATGCTGCTCCCGCACTCAACGTCGGGAGGACTGGCAGGGG-4800
 GAAGCCAGGCATCTACAGATTGTTGGCACCGGGGAGCGCCCTCCGGCATGTTCCGACTC
 GTCCGTCTCTGTGAGTGCTATGACGCAGGCTGTGCTGGTATGAGCTCACGCCCGCCGA
 GACTACAGTTAGGCTACGAGCGTACATGAACACCCCGGGCTTCCGTGTGCCAGGACCA
 TCTTGAATTTGGGAGGGCGTCTTTACAGGCCCTCACTCATATAGATGCCACTTTCATC
 CCAGACAAGCAGAGTGGGGAGAACCTTCTTACCTGCTAGCGTACCAAGCCACCGTGTG-5100
 CGTAGGGCTCAAGCCCTCCCCATCGTGGACCAAGATGTGGAAGTGTGATTGCTGCCCT
 CAAGCCACCCCTCCATGGGCCAACACCCCTGCTATACAGACTGGGCGCTGTTCAGAATGA
 AATCACCCCTGACGCCACCCAGTCAACCAATACATCATGACATGTCGCGCCGACCTGGA
 GGTCGTACGAGCACCTGGGTGCTGCTGGGGGCTCTGGCTGCTTTGGCCGCGTATTG
 CCTGTCAACAGGCTGCGTGGTCAATAGTGGGAGGTCGCTTGTCCGGGAGCCGGCAAT-5400
 CATACCTGACAGGGAAGTCTCTACCGAGAGTTCGATGAGATGGAAGAGTCTCTCAGCA
 CTTACCGTACATCGAGCAAGGATGATGCTCGCGAGCAGTTCAAGCAGAAAGCCCTCGG
 CCTCCTGCAGACCGGCTCCCGTCAGGCAGAGGTTATCGCCCTGCTGCCAGACCAACTG
 GCAAAAACCTCGAGACCTTCTGGGCGAAGCATATGTGGAACCTTCATCAGTGGGATACAATA
 CTGGCGGGCTTGTCACGCTGCCGTGTAACCCCGCCATTGCTTCAATTGATGGCTTTTAC-5700
 AGCTGCTGTACACGCCCACTAACCTAGCCAAACCTCCTCTTCAACATATTGGGGG
 GTGGGTGGCTGCCAGCTCGCCGCCCGGTCGCTACTGCTTGTGGCGCTGGCTT
 AGCTGGCGCGCCATCGGCAGTGTGGACTGGGAAGGTCCTCATAGACATCCTTGCAGG
 GTATGGCGCGGCGTGGCGGAGCTCTGTGGCATTCAGATCATGAGCGGTGAGTCCC
 CTCCACGGAGGACCTGGTCAATCTACTGCCCGCCATCCTCTCGCCCGAGCCCTCGTAGT-6000
 CGCGGTGCTGTGCAGCAATACTGCGCGGCACTGTTGGCCCGGGAACCATGTTCCCGCACGCACTA
 GTGGATGAACCGGCTGATAGCTTCCGCTCCCGGGGAGGAGTGTTCCTCCCGACGCACTA
 CGTGCCGAGAGCGATGAGCTGCCGCGTCACTGCCATACCTCAGCAGCCCTCACTGTAACT
 CCAGCTCCTGAGGCGACTGCACCGAGTGATAGCTCGGAGTGTACCACTCCATGCTCCGG
 TTCTCTGGCTAAGGGACATCTGGGACTGGATATCGGAGGTGTTGAGCGCACTTAAAGACCTG-6300
 GCTAAAAGCTAAGCTCATGCCACAGCTGCCCTGGGATCCCCCTTTGTGTCTGCTCCAGCGCGG

WO 93/00365

PCT/US92/05388

8 / 23

FIG.2-5

GTATAAGGGGTCTGGCGAGTGGACGGCATCATGCACACTCGCTGCCACTGTGTGGAGCTGA
 GATCACTGGACATGTCAAAAACGGGACGATGAGGATCGTGGTCTTAGGACCTGCAGGAA
 CATGTGAGTGGACCTTCCCATTAATGCTACACACGGGCCCTGTACCCCTTCC
 TGGCCGAACATACAGTTCGCGCTATGGAGGTGTCTGCAGAGGAATATGTGGAGATAAG-6600
 GCAGGTGGGGACTTCCACTACGTACGGGTATGACTACTGACAAATCTCAATGCCCTG
 CCAGGTCCCATCGCCGAAATTTTACAGAAATGGACGGGTGCGCTACATAGGTTGC
 GCGCCCTGCAAGCCCTTGCTGCGGAGGAGGATCATTCAGAGTAGGACTCCACGAATA
 CCGGTAGGTGCAATTAACCTTGCAGCCCGAACGGACGTGGCCGTGTGACGTCCAT
 GCTCACTGATCCCTCCCATATAACAGCAGAGGCGCGGCGGAGGTTGGCAGGGGATC-6900
 ACCCCCTCTGTGGCCAGCTCCTCGGCTAGCCAGCTATCCGCTCCATCTCTCAAGGCAAC
 TTGCACCGCTAACCATGACTCCCTGATGCTGAGCTCATAGAGGCCAACCTCTATGGAG
 GCAGGAGATGGCGGCAACATCACAGGTTGAGTCAGAAACAAAGTGGTGTCTTGA
 CTCCTTCGATCCGCTTGTGGCGGAGGAGGACGAGCGGAGATCTCCGTACCCGCAGAAAT
 CCTGCGGAAGTCTCGGAGATTCGCCAGGCCCTGCCCTTGGCGCGGCCGACTATAA-7200
 CCCCCGCTAGTGGAGACGTGGAAAAGCCCGACTACGAACCACTGTGGTCCATGGCTG
 TCCGCTTCCACCTCCAAAGTCCCTCTGCTGCTCCGCTCGGAAGAACGGACGGTGGT
 CCTCACTGAATCAACCTATCTACTGCCCTTGGCCGAGCTCGCCACCAGAACTTTGGCAG
 CTCCCTCAACTCCGGCATTACGGCGCAATACGACAAACATCCTCTGAGCCCGCCCTTC
 TGGCTGCCCCCGACTCCGACGCTGAGTCTTATCTCTCCATGCCCTGGAGGGGA-7500
 GCCTGGGATCCGGATCTTAGCGACGGGTCACTGGTCAACGGTCAGTAGTGAAGCCAAAGC
 GGAGGATGTCGTGCTCAATGCTTACTCTTGGACAGGCGCACTCGTCAACCCCGTG
 CGCCGGGAAGAACAGAACTGCCCATCAATGCACATAAGCAACTCGTTGCTACGTACCCA
 CAATTGGTGTATTCCACCACCTCACGCGAGTGTGTCCTGTCCTGAGGAGGTTAAAGCAGC-7800
 TGACAGACTGCAAGTCTGACAGCCATTACAGGACGTACTCAAGGAGGTTAAAGCAGC
 GGCGTCAAAAGTGAAGGCTAATCTGCTATCCGTAGAGGAAGCTTGCAGCTGACGCCCC
 AACTCAGCCAAATCCAAAGTTTGGTATGGGCAAAAGACGTCCTGTCATGCCAGAA
 GGCCGTAACCCACATCAACTCCGTGTGGAAGACCTTCTGGAAGACAAATGTAAACCCAAAT
 AGACACTACCATCATGGCTAAGAACGAGGTTTCTGCGTTTCAGCCCTGAGAAAGGGGTCTG

WO 93/00365

PCT/US92/05388

9 / 23

FIG.2-6

TAAGCCAGCTCGTCTCATCGTGTTCCTCCCGATCTGGGCGTGCGGCGTGTGCGAAAGATGGC-8100
 TTTGTACGACGTGGTTACAAAGCTCCCTTGGCCGTGATGGGAAGCTCCTACGGATTCCA
 ATACTCACGAGACAGCGGTTGAATTCCTCGTGCAAGCGTGAAGTCCAGAAACCC
 AATGGGTTCTCGTATGATACCCGCTGCTTGTACTCCACAGTCACTGAGAGCGACATCCG
 TACGGAGGAGGCAATCTACCAATGTTGTGACCTCGACCCCAAGCCCGTGGCCATCAA
 GTCCCTCACCGAGAGGCTTTATGTTGGGGCCCTCTTACCAATCAAGGGGAGAACTG-8400
 CGGCTATCGCAGGTCCGCGGAGCGGCTACTGACAACTAGCTGTGGTAACACCCCTCAC
 TTGCTACATCAAGGCCCGGCAGCCTGTCGAGCCGAGGGCTCCAGGACTGCACCATGCT
 CGTGTGGCGACGACTTAGTCTTATCTGTGAAGCGGGGGTCCAGGAGGACGCGGC
 GAGCCTGAGAGCCTTCACGGAGGCTATGACAGGTACTCCGCCCCCTCGGGGACCCCC
 ACAACCAGAAATACGACTTGGAGCTCATACATGCTCTCTCAACGTGTAGTGCCTCA-8700
 CGACGGCGCTGGAAGAGGGTCTACTACCTCACCCGTGACCTACACCCCTCGCGAG
 AGCTGCGTGGGAGACAGCAAGACACACTCCAGTCAATTCCTGGCTAGGCAACATAATCAT
 GTTTGCCCCCACACTGTGGCGAGGATGATAGTATGATACCCATTTCTTAGCGTCCCTAT
 AGCCAGGACAGCTTGAACAGGCCCTCGATTGCGAGATCTACGGGGCTGCTACTCCAT
 AGAACCACTTGATCTACCTCCAATCATTCAAAGACTCCATGGCCTCAGCGCAATTTCACT-9000
 CCACAGTTACTCTCCAGGTGAAATTAATAGGGTGGCGCATGCCTCAGAAAACTTGGGGT

 ACCGCCCTTGGAGACTTGGAGACACCGGGCCCGAGCGTCCGCGCTAGGCTTCTGGCCAG
 AGGAGCGAGGCTGCCATATGTGGCAAGTACCTCTTCAACTGGGCAGTAAGAACAAAGCT
 CAAACTCACTCCAATAGCGCGCTGGCCAGCTGGACTTGTCCGGCTGGTTACGGCTGG
 CTACAGCGGGGAGACATTTATCACAGCGTGTCTCATGCCGCGCCCGCTGGATCTGGTT-9300
 TTGCCCTACTCCTGCTGTGTCAGGGGTAGGCATCTACCTCCTCCCCAACCGATGAAGGTT
 GGGGTAAACACTCCGGCCT-----3' terminus

Some clonal heterogeneities producing amino acid substitutions are shown. There are many other "silent" mutations(not shown).

WO 93/00365

PCT/US92/05388

10 / 23

FIG.3-1 Human 23

1 GlyPheAlaAspLeuMetGlyTyrIleProLeuValGlyAlaProLeuGlyGlyArgAla
 GGTTCCGCCGACCTCATGGGTACATACCGCTCGTCGGCGCCCTCTTGGAGGCCGTGCC
 61 ArgAlaLeuAlaHisGlyValArgValLeuGluAspGlyValAsnTyrAlaThrGlyAsn
 AGGCCCTGGCGCACGGCTCCGGGTTTGGAGACGGCGTGAACATATGCAACAGGGAAC
 CG A
 LeuProGlyCysSerPheSerIlePheLeuLeuAlaLeuLeuSerCysLeuThrValPro
 121 CTTCCTGGTTGCTCCTTTTCTATCTTCCTTCTGGCCCTACTCTCTTGCCCTGACCGTGCCC
 GA
 AlaSerAlaTyrGlnValArgAsnSerThrGlyLeuTyrHisValThrAsnAspCysPro
 181 GCTTCAGCCTACCAAGTGGCAACTCTACGGGGCTTTACCATGTCACCAATGATGCCCCT
 AsnSerSerIleValTyrGluAlaAlaAspAlaIleLeuHisAlaProGlyCysValPro
 241 AACTCGAGTATTGTACGAGGCGCGCGATGCCATCCTGCACGCTCCGGGTGTGCCCT
 T C
 CysValArgGluAspAsnValSerArgCysTrpValAlaValThrProThrValAlaThr
 301 TCGGTTCCGAGGATAACGTCTCGAGATGTTGGGTGGGTGACCCACGGTGCCACC
 G T
 LysAspGlyLysLeuProThrThrGlnLeuArgArgHisIleAspLeuLeuValGlySer
 361 AAGGACGGCAAACTCCCAACGACGCTTCGACGTACATCGATCTGCTTGTCGGGAGC
 C C
 AlaThrLeuCysSerAlaLeuTyrValGlyAspLeuCysGlySerIlePheLeuValGly
 421 GCCACCCCTCTGCTCGGCCCTCTACGTGGGGACCTTTGCGGGTCCATCTTTCTTGTTCGGT
 T
 GlnLeuPheThrPheSerProArgArgHisTrpThrThrGlnAspCysAsnCysSerIle
 481 CAACTGTTTACCTTCTCTCCAGCGCCCACTGGACGACGACGACTGCAACTGTTCTATC
 C

WO 93/00365

PCT/US92/05388

11 / 23

FIG.3-2

```

541 TyrProGlyHisIleThrGlyHisArgMetAlaTrpAspMetMetMetAsnTrpSerPro
    TATCCCGGCATATAACGGGTCACCGCATGGCATGGATATGATGATGAATGAACTGGTCCCCCT
        G
601 ThrAlaAlaLeuValValAlaGlnLeuLeuArgIleProGlnAlaIleLeuAspMetIle
    ACGCGGCATTGGTAGCTAGCTCAGCTGCTCCGGATCCCAAGCCATCTTGGACATGATC
        G   AG
661 AlaGlyAlaHisTrpGlyValLeuAlaGlyMetAlaTyrPheSerMetValGlyAsnTrp
    GCTGGTGCTCACTGGGAGTCCCTGGCGGCATGGCGTATTTCTCCATGGTGGGAACTGG
        G
721 AlaLysValLeuValLeuLeuLeuPheAlaGlyValAlaAspAlaGluThrHisArgThr
    GCGAAGGTCTCTGGTAGTGCTGCTTCTATTGCGCGTCTGACGCGGAAACCCACCGTACC
        G
781 GlyGlySerAlaAlaArgSerThrAlaGlyValAlaSerLeuPheThrProGlyAlaArg
    GGGGGAAGTGCCGCCCGCAGCAGCGGTGGAGTTGCTAGTCTCTTCACACCCAGGCGCTAGG
    C      T      A
841 GlnAsnIleGlnLeuIleAsnThrAsnGlySerTrpHisIleAsnSerThrAlaLeuAsn
    CAGAACATCCAGCTGATCAACACCAACGCGCAGTTGGCACATCAATAGTACGGCCCTTGAAC
        AT
901 CysAsnAspSerLeuThrThrGlyTrpLeuAlaGlyLeuPheTyrHisHisLysPheAsn
    TGCAATGACAGCCTTACCACCGGCTGGTTAGCGGGCTTTTCTATCACCATAAATTCAAC
    A      A
961 SerSerGlyCysProGluArgLeuAlaSerCysArgProLeuThrAspPheAlaGln
    TCTTCAGGCTGTCCCGAGAGGTTGGCCAGCTGCCGACCCCTCACCGATTTTGCCCCAGG
    G      A

```

WO 93/00365

PCT/US92/05388

12 / 23

FIG.4-1 Human 27

GlyPheAlaAspLeuMetGlyTyrIleProLeuValGlyAlaProLeuGlyGlyAlaAla
GGCTTCGCCGACCTCATGGGTACATTCCGCTCGTCGGCTCCTCTTGGGGCGCTGCC
1
61 ArgAlaLeuAlaHisGlyValArgValLeuGluAspGlyValAsnTyrAlaThrGlyAsn
AGGGCCCTGGCGCATGGCGTCCGGTCTCTGGAAGACGGCGTGAACATATGCAACAGGAAC
121 LeuProGlyCysSerPheSerIlePheLeuLeuAlaLeuLeuSerCysLeuThrValPro
CTTCCCTGGTTGCTCTTTCTCTATCTTCTCTGGCTCTGCTCTCTTGCCTGACCGTGCCC
181 AlaSerAlaTyrGlnValArgAsnSerSerGlyIleTyrHisValThrAsnAspCysPro
GCATCGGCCTACCAAGTACGCAACTCCTCGGGCATTTACCATGTCAACCAATGATTGCCCT
241 AsnSerSerIleValTyrGluThrAlaAspThrIleLeuHisSerProGlyCysValPro
AATTCGAGTATTGTGTACGAGACGGCCGACACCATCTACACTCTCCGGGTGTGTCCCT
C
301 CysValArgGluGlyAsnAlaSerLysCysTrpValProValAlaProThrValAlaThr
TGCGTTCGCGAGGGTAACGCCCTCGAAATGTTGGTGCCGGTAGCCCCCACAGTGGCCACC
G
361 ArgAspGlyAsnLeuProAlaThrGlnLeuArgArgHisIleAspLeuLeuValGlySer
AGGGACGGCAACCTCCCCGCAACGCAGCTTCGACGTCACATCGATCTGCTTGTCTGGGAGT
G
421 AlaThrLeuCysSerAlaLeuTyrValGlyAspLeuCysGlySerValPheLeuValGly
GCCACCCCTTGTCTCGGCCCTCTATGTGGGGACTTGTGCGGGTCTGTCTTTCTTGTCTGGT
C
481 GlnLeuPheThrPheSerProArgArgHisTrpThrThrGlnAspCysAsnCysSerIle
CAACTGTTCACCTTTCTCCCCCAGGCGCCACTGGACAACGCAAGATTGCAACTGCTCTATC
A

WO 93/00365

PCT/US92/05388

13 / 23

FIG. 4-2

541 TyrProGlyHisIleThrGlyHisArgMetAlaTrpAspMetMetMetAsnTrpSerPro
TACCCCGGCATATAACGGACACCGCATGGCATGGGATATGATGATGAACTGGTCCCCCT

601 ThrAlaAlaLeuValMetAlaGlnLeuLeuArgIleProGlnAlaIleLeuAspMetIle
ACAGCAGCGCTGGTAATGGCTCAGCTGCTCAGGATCCCGAAGCCATCTTGGACATGATC
G

661 AlaGlyAlaHisTrpGlyValLeuAlaGlyIleAlaTyrPheSerMetValGlyAsnTrp
GCTGGTGCTCACTGGGAGTCTCTAGCGGCATAGCGTATTCTCCATGGTGGGAACTGG

721 AlaLysValLeuValValLeuLeuLeuPheAlaGlyValAspAlaThrThrTyrThrThr
GCGAAGTCTCTGGTGGTGCTGTGTGCTGTTGCGCGGTCGATGCGACAACCTATACCACC

781 GlyGlyAsnAlaAlaArgThrThrGlnAlaLeuThrSerPhePheSerProGlyAlaLys
GGGGGAATGCTGCCAGGACCACGCGCGCTCACCAGTTTTCAGCCCAGGCGCCAAG

841 GlnAspIleGlnLeuIleAsnThrAsnGlySerTrpHisIleAsnArgThrAlaLeuAsn
CAGGATATCCAGCTGATCAACACCAACGCGCAGTTGGCACATCAATCGCACGGCCTTGAAAC
G

901 CysAsnAlaSerLeuAspThrGlyTrpValAlaGlyLeuPheTyrTyrHisLysPheAsn
TGTAATCGGAGCCTCGACACTGGCTGGGTAGCGGGCTCTTCTATTACCACAAATTCAAC
T

961 SerSerGlyCysProGluArgMetAlaSerCysArgProLeuAlaAspPheAspGln
TCCTCAGGCTGCCCCGAGAGGATGGCCAGCTGTAGGCCCCCTTGCCGATTTCGACCAGG
C

WO 93/00365

PCT/US92/05388

14 / 23

FIG.5-1

1. human 27 2. HCV 1 3. human 23

```

1 CGGCTTCGCCGACCTCATGGGGTACATtCCGCTCGTCGGCGctCCTCTTGGgGGCGCTGCCAGGGCCCTGGC
*****
1 CGGCTTCGCCGACCTCATGGGGTACATACCGCTCGTCGGCGCCCTCTTGGAGCGCTGCCAGGGCCCTGGC
*****
1 CGGCTTCGCCGACCTCATGGGGTACATACCGCTCGTCGGCGCCCTCTTGGAGCGcgTGCCAGGGCCCTGGC
*****
73 GCATGGCGTCCGGGTCTGGAAGACGGCGTGAACtATGCAACAGGGAACCTTCCTGGTGTCTCTTCTCTAT
*****
73 GCATGGCGTCCGGGTCTGGAAGACGGCGTGAACtATGCAACAGGGAACCTTCCTGGTGTCTCTTCTCTAT
***
73 GCACGGCGTCCGGGTtTGAAGACGGCGTGAACtATGCAACAGGGAACCTTCCTGGTGTCTCTTCTCTAT
*****
145 CTTCCTTCTGGCtCTGCTCTCTTGcCTGACcGTGCCCGCbTCGGCCTACCAAGTaCGCAACTCctCGGGcAt
*****
145 CTTCCTTCTGGCCCTGCTCTCTTGcCTGACtGTGCCCCGCTTCGGCCCTACCAAGTGCgCAACTCCACGGGGCT
*****
145 CTTCCTTCTGGCCCTaCTCTCTTGcCTGACcGTGCCCGCTTCaGCCTACCAAGTGCgCAACTCtACGGGGCT
*****
217 TTACCAtGTCACCAATGATtGCCCTAAtTCGAGTATtGTGTACGAGaCGGCCGACaCCATCCTaCACtCTCC
*****
217 TTACCAcGTCACCAATGATtGCCCTAACTCGAGTATtGTGTACGAGCGGCCGATGCCATCCTGCACaCTCC
*****
217 TTACCAtGTCACCAATGATtGCCCTAACTCGAGTATtGTGTACGAGCGGCCGATGCCATCCTGCACgCTCC
*****
289 GGGGTGtGTCCCTTGCGTTCGGGAGGGtAACGCCCTCGAaaTGTTGGGTGcCGgTagCCCCcACaGTGGCCAC
*****
289 GGGGTGcGTCCCTTGCGTTCGtGAGGGcAACGCCCTCGAGgTGTTGGGTGGCGaTGACCCCTtACGGTGGCCAC
*****
289 GGGGTGtGTCCCTTGCGTTCGGcGAGGatAACGtCTCGAGaTGTTGGGTGGCGgTGACCCCCcACGGTGGCCAC
*****

```

WO 93/00365

PCT/US92/05388

361 CAGGGACGGCAACCTCCCGCaACGCAGCTTCGACGTCAATCGATCTGCTGTGTCGGAGtGCCACCCtTG

361 CAGGGAtGGCAAACTCCCGGgACGCAGCTTCGACGTCAATCGATCTGCTGTGTCGGAGCGCCACCCtCTG
**
361 CAaGGACGGCAAACTCCCCaCaACGCAGCTTCGACGTCAATCGATCTGCTGTGTCGGAGCGCCACCCtCTG

433 cTCGGCCCTCTAtGTGGGGACtTGTCGGGTCTGCTTTCTTGTCGGtCAACTGTTcACTtTTCtTcCCCCAG

433 tTCGGCCCTCTACGTGGGGACCTGTGCGGGTCTGCTTTCTTGTCGGcCAACTGTTcACTTCTCTCTCCAG

433 cTCGGCCCTCTACGTGGGGACCTtTGCGGGTccaTCTTTCTTGTCGGtCAACTGTTtACCTTCTCTCCAG

505 GCGCCACTGGACaACGCAAGaTTGCAAcTGCTCTATcCCCGGCCATATAACGGGaCACCgCATGGCATG

505 GCGCCACTGGACGACGAAGgTTGCAAtTGCTCTATcCCCGGCCATATAACGGGTcACCGCATGGCATG¹

505 GCGCCACTGGACGACGAgaCTGCAAcTGtTCTATCTATcCCCGGCCATATAACGGGTcACCGCATGGCATG²

577 GGATATGATGATGAACtGGTCCCTACagCaGCGcTGgTAATGCTCAGCTGCTCaGGATCCCGCAAGCCAT

577 GGATATGATGATGAACtGGTCCCTACGaCGGCGTTGgTAATGCTCAGCTGCTCCGGATCCCAAGCCAT

577 GGATATGATGATGAACtGGTCCCTACGgCGGCaTTGgTAgtAGCTCAGCTGCTCCGGATCCCAAGCCAT

649 CTtGGACATGATCGCTGGTCTCACTGGGGAGTCCtAgCGGGCATAGCGTAATTCTCCATGGTGGGAACtG

649 CTtGGACATGATCGCTGGTCTCACTGGGGAGTCCtGCGGGCATAGCGTAATTCTCCATGGTGGGAACtG

649 CTtGGACATGATCGCTGGTCTCACTGGGGAGTCCtGCGGGCATgGCCGTAATTCTCCATGGTGGGAACtG

721 GCGGAAGTCCtGGTgGTGCTgtTGCTgTTTGCCGGCTCGAtCGGacAAcCtAtacCACCgGGGgAaTGC

721 GCGGAAGTCCtGGTAGTGTGCTGTCTATTTGCCGGCTCGACGCGGAACCCACgtCACCGGGGGAAGTGC

721 GCGGAAGTCCtGGTAGTGTGCTtCTATTTGCCGGCTCGACGCGGAACCCACcgTACCgGGGGAAGTGC

WO 93/00365

PCT/US92/05388

16 / 23

793 tGcCagGACcacGcagGcgCTaccAGtTtTTCagcCCAGGCGCCAAGCAGgAtaTCCAGCTGATCAACAC
* * * * *
793 CGgCCaCActgtGtCTGGAtTTGtTAGcCTCcTCgCACcAGGCGCCAAGCAGAACgTCCAGCTGATCAACAC
* * * * *
793 CGcCCgCagcacGgCTGGAgTTGcTAGtCTCtTcCaCACcAGGCGCtAgGCAGAACaTCCAGCTGATCAACAC

865 CAACGGCAGTTGGCACA TCAATcGCACGGCCtTGAACtGtAATGcgAGCCTCgACACtGGCTGGgTaGCgGG
* * * * *
865 CAACGGCAGTTGGCACcTCAATAGCACGGCCcTGAACtGCAATGAtAGCCtCAACACCGGCTGGTTgGCaGG
* * * * *
865 CAACGGCAGTTGGCACaTCAATAGtACGGCCtTGAACtGCAATGAcAGCCtTAcCACCGGCTGGTTaGCgGG

937 GCTcTTCTATtACCACAAaTTCAACTCTTCAGGCTGcCCcGAGAGGaTgGCCAGCTGtagGCCCTTgCCGA
* * * * *
937 GCTTTCTATCACCACAAGTTCAACTCTTCAGGCTGTCCtGAGAGGcTaGCCAGCTGCCGACCCCTTACCGA
* * * * *
937 GCTTTCTATCACCAtAAaTTCAACTCTTCAGGCTGTCCcGAGAGGtTgGCCAGCTGCCGACCCCTcACCGA

1009 TTTGACCAGG
*** *****
1009 TTTGACCAGG
*** *****
1009 TTTGACCAGG

FIG.5-3

FIG.6

1	GFADLMGYIPLVGAPLGGARALAHGVRVLEdGVNYATGNLPGCSFSIFLLALLSCLTVPASAYQVRNSSGi	
1	GFADLMGYIPLVGAPLGGARALAHGVRVLEdGVNYATGNLPGCSFSIFLLALLSCLTVPASAYQVRNSTGL	
1	GFADLMGYIPLVGAPLGGrARALAHGVRVLEdGVNYATGNLPGCSFSIFLLALLSCLTVPASAYQVRNSTGL	
'3	YHVTNDCPNSSIVYEtADtILHsPGCVPCVREGNASKCWvpvaPTVATRdGnLPATQlRRHIDLLVGSATLC	
'3	YHVTNDCPNSSIVYEAADAiLHtPGCVPCVREGNASRCWvAmTPTVATRdGKLpATQlRRHIDLLVGSATLC	
'3	YHVTNDCPNSSIVYEAADAiLHaPGCVPCVREDnVSRcWvAvTPTVATkdGKLpTQlRRHIDLLVGSATLC	
5	SALyVGDLcGSvFLVGQLFTFSPrRHwTtQdCNCsIYPGHITGHRMAWdMMNWSPTaALVMAQLLRIPQAI	
5	SALyVGDLcGSvFLVGQLFTFSPrRHwTtQ9CNCsIYPGHITGHRMAWdMMNWSPTtALVMAQLLRIPQAI	
5	SALyVGDLcGSiFLVGQLFTFSPrRHwTtQdCNCsIYPGHITGHRMAWdMMNWSPTaALVvAQLLRIPQAI	
7	LDMIAGAhWGVLAGIAYFSMVGNWAKVLVLLLFAGVDAtTyttGGnAarTtqaltSffsPGAKQdiQLINT	
7	LDMIAGAhWGVLAGIAYFSMVGNWAKVLVLLLFAGVDAETHvTGGSaghtVsGfvSLlapGAKQNVQLINT	
7	LDMIAGAhWGVLAGmAYFSMVGNWAKVLVLLLFAGVDAETHrTGGSaArstaGvaSLftPGARQNiQLINT	
39	NGSWHiNrTALNCNaSLdTGWvAGLFYyHKFNSSGCPERmASCRPLaDFDQ	
39	NGSWHlNSTALNCNDSLnTGWLAGLFYHHKFNSSGCPERlASCRPLTDfDQ	
39	NGSWHiNSTALNCNDSLtTGWLAGLFYHHKFNSSGCPERlASCRPLTDfAq	
		1. human 27
		2. HCV 1
		3. human 23

WO 93/00365

PCT/US92/05388

CLUSTERED PAIR-WISE 'REGION' ALIGNMENT
in 'identity (no translation)' alphabet of:

FIG. 7 -- I

1. ssThorn#8.r (1-587)
2. sSEC1#2.r (1-587)
3. sSHCT18#7.r (1-587)
4. env1.hcv (1-1657)

GA
||
GA
||
GA
||

1
1
1


289 gggtaggggagtggtcctgtctccccgtggctctcgccctagctggggccacagacccccggcgtagg
3 ATTGCGCAATTGGGTAAGGTCAATCGATACCCCTTACGTGCGGCTTCGCCGACCTCATGGGGTACATACCGCTC
3 ATTGCGCAATTGGGTAAGGTCAATCGATACCCCTTACGTGCGGCTTCGCCGACCTCATGGGGTACATACCGCTC
3 ATTGCGCAATTGGGTAAGGTCAATCGATACCCCTTACGTGCGGCTTCGCCGACCTCATGGGGTACATACCGCTC
361 t-cgCGCAATTGGGTAAGGTCAATCGATACCCCTTACGTGCGGCTTCGCCGACCTCATGGGGTACATACCGCTC

18
23

75 GTCGGCGCCCCCTCTTGGgGGCGCTGCCAGGGCCCTGGCGCATGGCGTCCGGGTTCTTGGAAGACGGCGTGAAC
75 GTCGGCGCCCCCTCTTGGAGGCGCTGCCAGGGCCCTGGCGCATGGCGTCCGGGTTCTTGGAAGACGGCGTGAAC
75 GTCGGCGCCCCCTCTTGGAGGCGCTGCCAGGGCCCTGGCGCATGGCGTCCGGGTTCTTGGAAGACGGCGTGAAC
33 GTCGGCGCCCCCTCTTGGAGGCGCTGCCAGGGCCCTGGCGCATGGCGTCCGGGTTCTTGGAAGACGGCGTGAAC

PCT/US92/05388

147 TATGCAACAGGAAACCTTCCCTGGTTGCTCTTTCTCTcTCTTCCCTTCTGGCCcTGCTCTCTTGtctTGACcCGTG
147 TATGCAACAGGAAACCTTCCCTGGTTGCTCTTTCTtTATCTTCCCTTCTGGCCcTGCTCTCTTGCTtTGACTGTG
147 TATGC CAGGGAACCTTCCCTGGTTGCTCTTTCTCTATCTTCCCTTCTGGCCCTGCTCTCTTGCCcTGACTGTG
505 TATGCAACAGGAAACCTTCCCTGGTTGCTCTTTCTCTATCTTCCCTTCTGGCCCTGCTCTCTTGCTtTGACTGTG
219 CCCGCTTCAGCCTACCAAGTGCAGCAACTCCaCGGGGCTTTACCAATGTACCCAACGAATTGCCCCcAACTCGAGt
219 CCCGCTTCAGCCTACCAAGTGCAGCAACTCCcCGGGGCTTTACCAATGTACCCAATGATTGCCCTAACTCGAGc
219 CCCGCTTCAGCCcACCAAGTGCAGCAACTCCACGGGGCTTTACCAATGTACCCAATGATTGCCCCcAACTCGAGT¹⁹
377 CCCGCTTCgCCc+ACCAAGTGCAGCAACTCCACGGGGCTTTACCAcGTACCCAATGATTGCCCTAACTCGAGT²³
191 ATTGTGTACGAGGCGCGATGctATCCTGCACgCTCCGGGGTGTGTCCCTTGCGTTcGCGAGGGtAAACGCC
191 ATTGTGTACGAGGCGCGCGATGCCATCCTGCACACTCCGGGGTGTGTCCCTTGCGTTcACGAGGGCAACGTC
91 ATTGTaTACGAaGCGGCCGAcGCCATCCTGTCAcACTCCGGGGTGTGTCCCTTGCGTTcACGAGGGCAACGTC
49 ATTGTgTACGAgGCGGCCGAtGCCATCCTGTCAcACTCCGGGGTGTcGTCCCTTGCGTTcgtGAGGGCAACGCC
63 TCGAGGTGTtGGGTGGCGATGACCCCCACGGTGGCCgCCAGGGaCGGCAGACTCCCCACAACGCAGCTgCGA
63 TCGAGGTGTtGGGTGGCGATGACCCCCACGGTGGCCcACcAGGGgCGGCAAACTCCCCACAACGCAGCTTCGA
63 TCGAGGTGTtGGGTGGCGgTGACCCCCACGGTGGCCACcAGGGATGGCAAACTCCCCACAACGCAGCTTCGA
21 TCGAGGTGTtGGGTGGCGaTGACCCCCtACGGTGGCCACcAGGGATGGCAAACTCCCCgCgACGCAGCTTCGA



WO 93/00365

PCT/US92/05388

21 / 23

```

          10      20      30      40
EC10      GAATTCGGACGACGCAAGGTTGCAATTGCTCTATCTATCCCGGCCATAT
          X::::::::::::::::::::::::::::::::::::::::::::::::::::
HCV1      CTCTCCCAGGCGCCACTGGACGACGCAAGGTTGCAATTGCTCTATCTATCCCGGCCATAT
          550      560      570      580      590      600

          50      60      70      80      A      90      100
AACAGGTCACCGCATGGCATGGGATATGATGATGAACTGGTCCCCTACGACGGCGT TAGT
          ::::::::::::::::::::::::::::::::::::::::::::::::::::::
AACGGGTCACCGCATGGCATGGGATATGATGATGAACTGGTCCCCTACGACGGCGTGGT
          610      620      630      640      650      660

110      120      130      140      150      160
GGTAGCTCAGCTGCTCCGGATCCCACAAGCCATCTTGGACATGATCGCTGGTGCTCACTG
          : ::::::::::::::::::::::::::::::::::::::::::::::::::::::
AATGGCTCAGCTGCTCCGGATCCCACAAGCCATCTTGGACATGATCGCTGGTGCTCACTG
          670      680      690      700      710      720

170      180      190      200      210      220
GGGAGTCCTGGCGGGCATAGCGTATTTCTCCATGGTGGGGAAC TGGGCGAAGGTCTTGGC
          ::::::::::::::::::::::::::::::::::::::::::::::::::::::
GGGAGTCCTGGCGGGCATAGCGTATTTCTCCATGGTGGGGAAC TGGGCGAAGGTCTTGGT
          730      740      750      760      770      780

230      240      250      260      270      280
AGTGCTGCTGCTATTTGCCGGCGTCGACGCGGAAACCCACGTCACTGGGGGGGATCGCCGC
          ::::::::::::::::::::::::::::::::::::::::::::::::::::::
AGTGCTGCTGCTATTTGCCGGCGTCGACGCGGAAACCCACGTCACTGGGGGGGAAGTGCCGC
          790      800      810      820      830      840

290      300      310      320      330      340
CAAACTACGGCTAGCCTTACTGGTCTCTTCAATTTAGGTGCCAAGCAGAACATCCAGCT
          : : : : : : : : : : : : : : : : : : : : : : : : : : : :
CCACACTGTGTCTGGATTTGTTAGCCTCCTCGCACCAGGCGCCAAGCAGAACGTCCAGCT
          850      860      870      880      890      900

350      360      370      380      390      400
GATCAACACCAACGGCAGTTGGCACATCAACAGGACGGCCTTGA ACTGCAATGATAGCCT
          ::::::::::::::::::::::::::::::::::::::::::::::::::::::
GATCAACACCAACGGCAGTTGGCACCTCAATAGCACGGCCCTGA ACTGCAATGATAGCCT
          910      920      930      940      950      960

410      420
CAACACCGGCTGGAATTC
          ::::::::::::::X
CAACACCGGCTGGTTGGCAGGGCTTTTCTATCACCACAAGTTCAACTCTTCAGGCTGTCC
          970      980      990      1000      1010      1020

```

FIG.8

WO 93/00365

PCT/US92/05388

22 / 23

AA #117-308 (putative envelope region) FIG.9

- | | |
|-----------------------|--------------------|
| 1) HCT #18 (USA) | 3 clones sequenced |
| 2) JH23 (USA) | ? |
| 3) JH 27 (USA) | ? |
| 4) PBL-Th (USA) | 2 clones sequenced |
| 5) EC1 (Italy) | 3 clones sequenced |
| 6) HCV-1 (chimpanzee) | multiple |

C/M ← T → S

1) (P)

2)

3)

4)

5)

6) RNLGKVIDTLTCGFADLMGYIPLVGAPLGGAARALAHGVRVLEDGVNYATGNL

1) H

2)

3) S T T

4) L

5) (F) S

6) PGCSFSIFLLALLSCLTVPASAYQVRNSTGLYHVTNDCPNSSIVYEADAILH

1) Y (H) V V T

2) A D V V K T

3) S PVA N

4) A A R T

5) H V T

6) TPGCVPCVREGNASRCWVAMTPTVATRDGKLPATQLRRHIDLLVGSATLCS

1)

2) I D

3) D

4)

5) I

6) ALYVGDLGSGVFLVGQLFTFSPRRHWTTQGCNCSI

SUMMARY: "S" AA117-308 (93%)

HCT#18, PBL-Th, EC1(Italy) have 97% hom logy with HCV-1

JH23 and JH 27 have 96% and 95% homology with HCV-1, respectively

WO 93/00365

PCT/US92/05388

23 / 23

AA#300-438 (C-terminal region of the putative envelope region and amino ~1/3 of NSI)

- 1) JH23 ?
 2) JH27 ?
 3) Japanese isolate (T. Miyamura) ?
 4) EC10 (Italy) 2 clones sequenced
 (one nt difference, which did not
 result in an amino acid change)
 multiple

5) HCV-1 (chimpanzee)

S ← → NSI

1) D

A V

2) D

A

3)

V S

VM V

4)

5) TTQGCNCSIYPGHITGHRMAWDMMMNSPTTALVMAQLLRIPQAILDMIAGA

1) M

R

ARSTA VA

2)

T YT

N AR TQALT F

3) L Y

I M

GH R

VQ VT TLT

4)

A

I AK TASLTA

5) HWGVLAGIAYFSMVGNWAKVLVLLLFAGVDAETHVTGGSAGHTVSGFVSL

1) FS R I I T V

2) FT DI I R A D

3) FR S K I V I R Q F

4) FNL I I R N

5) LAPGAKQNVQLINTNGSWHLNSTALNCNDSLNTGWL

SUMMARY: NS 1 AA 330-660

"Isolate"	%Homology (AA330-438)	%Homology (AA383-405)
JH23	83	57
JH27	80	39
Japanese	73	48
EC10 (Italy)	84	48

FIG. 10